



Habitat and Ecosystem Report

Terminos Lagoon

Fisheries Improvement Project Mexico Yucatan Peninsula blue crab-dipnet/pot/trap



September 2023

CeDePesca

Contenido

Summary	3
Acronyms	3
Introduction.....	4
Laguna de Términos	5
Tides.....	5
Sediments.....	6
Winds.....	6
Climatic seasons	6
Water physico chemical variables	6
Submerged Aquatic Vegetation (SAV)	10
Seagrasses and macroalgae.....	11
Macroalgae.....	13
Mangrove.....	13
Fauna	14
Benthos.....	15
Ichtyc community.....	15
Birds, reptiles and mammals	17
Socio-environmental characteristics.....	18
Annex 1. Systematic list of species captured in Laguna de Terminos, period 2016-2017 (taken from Irola, et al., 2021).....	29
Annex 2. Fish species richness in the Terminos Lagoon: An occurrence data compilation of four sampling campaigns along a multidecadal series. From Paz-Ríos, et al., 2021.....	32
Annex III. List of species of phytobenthos in Laguna de Términos (Ortega, 1995) .	35

Summary

This report includes a brief description of the bio-physical, biological and social processes that had taken place in Terminos Lagoon, as well as the latest information generated for this area. It was recognized in 1994 as a Flora and Fauna Protection Area. It maintains numerous fisheries, one of which is that of blue crab

In the last 20 years it has been influenced by anthropic and natural impacts, among which the hydraulic management in the basin and global warming stand out, which modifies the physicochemical characteristics of the water (temperature, salinity and pH) and with it changes in the regionalization (Santos Santoyo et al., 2021) towards an homogenization of the habitat as salinity values have improved (Ramos-Miranda, et al., 2015) and a reduction of its depth.

This lagoon supports 117 species of fish, 69 of macro crustaceans (Raz, 2010), and 172 of mollusks (Reguero et al., 2010) among others. Of the species of fish and crustaceans registered for the Terminos Lagoon, 43 are commercially exploited (Ramos Miranda, 2000; Ramos Miranda et al., 2005a) (Ramos Miranda et al., 2015a), among them, the blue crab *Callinectes sapidus*.

There are four economic, social and cultural processes in the region that had been established which are linked to extensive production: They are extraction of "Palo de Tinte", extraction of Chicle tree, intensive shrimp exploitation and extraction of hydrocarbons.


Acronyms

Acronym	Description
SAV	Submerged Aquatic Vegetation
FIP	Fisheries Improvement Project
INEGI	National Institute of Statistics and Geography
SEMARNAP	Environment, Natural Resources and Fisheries Secretariat
INE	Ecology National Institute
m	meters
psu	Practical salinity unit

Introduction

Terminos Lagoon is an area characterized by its biodiversity. It was recognized in 1994 as a Flora and Fauna Protection. It maintains numerous fisheries, one of which is that of blue crab.

The fishing grounds of the FIP suppliers are located in Sabancuy and towards Isla Aguada, both located north of the Terminos Lagoon .



In 2018 many fishermen in the area transferred their fishing effort to other more profitable fisheries such as fish and shrimp. This caused difficulties in obtaining the blue crab and with this, the strategic decision was made to increase the number of suppliers and expand the supply area in order, to ensure the same volume. Due to the above, product is currently purchased from Atasta and Puerto Rico and sometimes it is purchased from marketers who bring product from Nuevo Campechito and Sánchez Magallanes, Tabasco, although in small quantities. The largest amount of blue crab comes from the northern part of the Laguna de Terminos.

Terminos lagoon besides having an ecological importance it has great social and economic relevance, as an area of intense economic development with population growth and industrial or extractive activities that have led to the modification of the ecosystem and habitats (Flores-Hernández, et al., 2021)

For example, the urban population of Ciudad del Carmen increased almost 29 times from 1940 to 2010 (INEGI, 2016). The oil industry developed, with impacts on fauna (Gold-Bouchot et al., 1995) and agriculture with the use of pesticides and fertilizers (Rendon-von Osten et al., 2005). On the other hand, natural phenomena such as hurricanes have modified the composition of channels. The sandy islands inside the

lagoon have spread, modifying the movement of the water and consequently the salinity. (Flores-Hernández, et al., 2021)

In the last 20 years it has been influenced by anthropic and natural impacts, among which the hydraulic management in the basin and global warming stand out, which modifies the physicochemical characteristics of the water (temperature, salinity and pH) and with it changes in the regionalization. (Santos Santoyo et al., 2021)

Laguna de Términos

Terminos Lagoon is located in the southwestern area of the Gulf of Mexico (Campeche State, Mexico). This is one of the largest lagoons in Mexico with a surface of 1660 km², forming part of the Tropical North-Western Atlantic's marine ecoregion 68 (Southern Gulf of Mexico). The lagoon is very shallow with a mean depth of 3.5 m. (Villéger, et al., 2010) Terminos Lagoon is actually an estuarine ecosystem as it is strongly influenced by freshwater discharges from three streams located on its southern part (respectively, from west to east: Palizada River, Chumpan River, and Candelaria River). Throughout the years, several studies show that the lagoon presents important changes associated with an environmental modification, especially salinity.

The lagoon is delimited by Carmen Island (30 km long and 2.5 km wide) which separates it from the Gulf of Mexico and thus water exchanges with the sea take place through two inlets, one on its northeastern part (Puerto Real) and the other one on the northwestern part (Carmen) (Villéger, et al., 2010) with depth of 18 m and 14 m respectively (Romo-Ríos, 2013), receiving freshwater from three streams on its southern part: Palizada River, Chumpan River and Candelaria River (Ramos-Miranda, et al., 2005a).. The mouth of Carmen, which is 3.9 km wide, has a fluvial influence derived from the Palizada River, which causes little transparency of the water due to the terrigenous material in suspension. (Borges Souza, 2004, in Romo-Ríos, 2013).

Water circulation in the lagoon generally follows a clockwise direction (David and Kjerfve 1998), with seawater going inside the lagoon through the Puerto Real inlet, mixing with freshwater inputs near the stream mouth, with the resulting brackish water going outside the lagoon through the Carmen inlet (Villéger, et al., 2010)

Tides

The tide in Terminos Lagoon is of a mixed type (diurnal and semi-diurnal) being the diurnal components the dominant ones; the range between mean high tide and mean low tide in Isla del Carmen is 0.43 m and the average high tide is 0.40 m and -0.70 m. Spring tide and neap tide have an approximate period of 14 days (SEMARNAP-INE, 1997 in Romo-Ríos, 2013).

Sediments

Termnos Lagoon's sediments are of biogenic and terrigenous origin, presenting spatio-temporal variations associated with the southwest-northeast circulation pattern present in the system, meteorological events, river discharges and the anthropogenic activity of the region. In a comparison of the composition and distribution of the sediments between results from 1963 and 2001-2002, it only presented significant differences in the Central Basin (Borges-Souza, 2004). The main sources of terrigenous sediments are the basins of the Chumpán and Palizada rivers (through the Usumacinta river on the Gulf of Mexico slope); the sources of carbonate sediment contribution are from the Gulf of Mexico through the Puerto Real Mouth and Mouth of Carmen, which are transported by tidal currents (Borges-Souza, 2004).

Winds

There are two dominant wind systems (Yáñez-Arancibia & Day, 1982):

a) northwest (October-March) with average speeds greater than 8 m/s, and b) north-northeast and east-southeast (March-September), with an average speed that varies between 4 and 6 m/s.

Climatic seasons

There are three climatic seasons in the area: dry, rainy and wind/winter season ("norte") seasons. The rainy season goes from June to October with a precipitation of 100 mm/month. Starting in November, with a slight decrease in precipitation, the wind/winter season goes until January. From February to May, the minimum precipitation values are recorded corresponding to the dry season (Yáñez Arancibia, et al. 1982).

The contribution of sediment to the lagoon derives from the Palizada rivers (discharge of 238,126 m³/s), flows into the Vapor and San Francisco lagoons through a small mouth of 7 m wide and 15 deep (Herrera Silveria et al., 2002), the Candelaria River with a discharge of 35.09 m³/s that flows into the Panlau Lagoon and connects to the Terminos Lagoon through a 40 m wide mouth. Finally, the Chumpán and Mamantel rivers with a discharge of 1.67 m³/s (Borges-Souza, 2004).

Water physico chemical variables

Throughout the years various studies have been carried out that describe the main physical and chemical variables of water: temperature, salinity, transparency and depth (Yáñez-Arancibia et al., 1982; Flores Hernández et al., 2001; Herrera-Silveira (2002) and Villalobos Zapata (2002), Sosa López et al. (2005)). these studies reflect the salinity and transparency gradients in the region and the fact that it is a dynamic environment that show from five to seven "habitats", whose borders are apparently environmentally changing (taken from Ramos et. Al, 2015a).

However, in the last 20 years it has been influenced by anthropic and natural impacts, which modifies the physicochemical characteristics of the water (temperature, salinity and pH) and changes in diversity species, being decisive in changes in regionalization

Santos, et.al. (2021) after an analysis of the spatio-temporal relationship of the physicochemical characteristics presented a new regionalization for Terminos Lagoon. Their results indicate for rainy season higher temperature values (30.59°C), higher salinity in dry (31.78 UPS), and higher pH in the wind/winter season (7.88). In the spatial distribution, the warmest areas were shown in the Candelaria River, the most saline areas by Isla de Carmen and the highest pH ranges near Pom-Atasta. 4 new zones are represented, which are influenced by fresh water discharges and the intrusion of salt water that is carried within the site. (Fig. 1)(Santos-Santoyo, et.al.,2021).





Fig 1. Map illustrating the 4 hydrological zones defined in Terminos Lagoon after Santos, et al., 2021.(a) (Illustration from Santos, et al., 2021) and Villéger et al, 2010 (b) (illustration from Ramos, et.al., 2015a)

The pH varies from 7.04 to 8.5 depending on the climatic season (dry, rainy and northerly), with higher pH during the wind/winter season (Santos, et.al., 2021).







Figure 2. Spatial variability of pH by climatic seasons (dry, rainy and wind/winter) after Santos et.al., 2021. (Illustration taken from Santos, et al., 2021)






Figure 3. Spatial variability of salinity by climatic seasons (dry, rainy and wind/winter), after Santos et.al., 2021. (Illustration taken from Santos, et al., 2021)

Villeger et.al. (2010) comparisons between 1980-1981 and 1998-1999 surveys showed that the four zones established by Yañez Arancibia (1982) experienced severe modifications in their environmental conditions (Table 1). Depth and transparency were globally decreasing, particularly in Zones 1, 2, and 3, while salinity increased Villéger et al., 2010.

Table I. Environmental conditions in four zones of the Terminos Lagoon (Villeger, et al., 2010)

Zone	Depth (m)		Sechii (m)		Bottom salinity (psu)		P	Substrate
	1980-1981	1998-99	P	1980-1981	1998-99	P		
1	2.8 (22 %)	2.1 (45 %)	***	0.6 (35 %)	0.5 (50 %)	*	21.3 (41 %)	23.1 (47%)
2	2.6 (32 %)	1.6 (45 %)	***	1.1 (45 %)	0.9 (49 %)	**	28.5 (25 %)	31.7 (18 %)
3	2.5 (24 %)	1.3 (35 %)	***	0.9 (38 %)	0.6 (39 %)	***	22.4 (28 %)	25.9 (22 %)
4	3.9 (11 %)	3.9 (9 %)	NS	1.1 (36%)	1.3 (44 %)	NS	26 (25 %)	31.1 (17 %)

Zone 1 is located near the mouth of Carmen, and is influenced by the discharge of freshwater from the Palizada River, whose annual discharge is $> 4 \times 10^9 \text{ m}^3 \text{ year}^{-1}$ and has a wide range of salinity (4 to 35 ups), presents a muddy substrate with fine sands and clays. (Villeger et al 2010)

Zone 2 covers the internal coastline of Isla del Carmen, up to the mouth of Puerto Real, this area is under marine influence with average salinities of 28.5 ups the substrate varies between muddy next to the mangrove strip (*Rhizophora mangle*) and sandy in seagrass areas (*Thalassia testudinum*). (Villeger et al 2010)

Zone 3 in the southeastern part of the lagoon, presents shallow waters close to mangrove swathes, it is influenced by the Candelaria and Chumpán rivers, with silty-clayey sediments. (Villeger et al 2010)

Zone 4 is the central part of the lagoon, with an average depth of 3.9 m, considered as a transition zone between the marine influence and the influence of fresh water, with a salinity range of 15 to 36 ups and an average of 26 ups. (Villeger et al 2010)

The behavior of the observed salinity has been interpreted by Ramos Miranda et al. (2015) as a homogenization of the habitat, pointing out that the greater intrusion of seawater and the pattern of circulation in the lagoon ecosystem, have conditioned a reduction in the wide ranges of salinity observed in the 80's. (Flores Hernández, 2021)

Submerged Aquatic Vegetation (SAV)

SAV can be found all around the Laguna de Terminos. There is evident relation between salinity gradients, freshwater inputs and the distribution of SAVs. *Calla lily*, *Eichornia crassipes*, cover large extensions toward the higher portion of the Palizada-del Este System. Rush marsh fields (*Typha sp*), as well as fields of reeds (*Phragmites*) can be found in those systems, being most abundant in the Palizada-del Este System. Other types of vegetation submerged in waters with low salinity values are *Vallisneria americana*, *Myriophyllum exaltatum* and *Potamogeton illinoensis*. (Robadue, et.al. 2004)




Figure 4. Map depicting distribution and location of Submerged Aquatic Vegetation and Surrounding Inundation Influenced vegetation in Laguna de Terminos Experts during Conservation Area Planning in May, 2004. (Illustration and information by Robadue, et al., 2004)

Seagrasses and macroalgae

The distribution of seagrasses and macroalgae is regulated by environmental conditions and affinity to textural groups of sediment and have been demonstrated as playing a key role in maintenance of high productivity in the region. Seagrasses are abundant in clear water (Robadue, et.al. 2004). Seagrass communities in Terminos Lagoon are dominated by *Thalassia testudinum* with two other species, *Halodule wrightii* and *Syringodium filiforme* occurring occasionally (Robadue, et.al., 2004).

The seagrasses are distributed in sandy and muddy sediments mainly on the southern coast of Isla del Carmen, through the floodtide delta at Puerto Real and along the littoral zone of the eastern and southeastern lagoon shorelines (Raz Guzman & Barba-Macías, 2000; Robadue, et.al. 2004). These regions are characterized by the highest water transparencies, salinities and percent calcium carbonates in the sediments. *Halodule* occurs on the shallowest flats while *Thalassia* extends to depths of 3 m. Greatest standing crops, leaf lengths and widths of *Thalassia* occur at depths between 1.0 and 2.0 m. Densities decrease with depth while ratios of aboveground to belowground biomass increase (Robadue, et.al., 2004). While the macroalgae are located in sites with sandy sediment such as the central part of Terminos Lagoon (Raz Guzmán & Barba-Macias, 2000). Its productivity varies between 3 and 13 gr of dry weight/m²/ day, with the highest rates occurring during spring in the area of Mouth of Puerto Real (Moore & Wetzel, 1988).




Figure 5. Map depicting distribution and location of seagrass beds in the Laguna de Terminos. Experts during Conservation Area Planning in May, 2004. (Illustration and information by Robadue, et al., 2004)

Yáñez-Arancibia et al. (1983), report a loss of *T. testudinum* of 37.5% of the coverage in the southeastern zone of Terminos Lagoon. In 1990, in the internal coastline of Isla del Carmen there was 58% coverage, while in Mouth of Puerto Real (West-East zone 2) it was 40% and in the river discharge zone (South zone 3) of 12%. By the year 2000 it was reduced to 38%, 35% and 5% respectively (Figure 6) (Herrera-Silveira et al., 2011).




Figure 6. Distribution and coverage of *Thalassia testudinum* in Terminos Lagoon (after Herrera-Silveira, et al., 2011)

For 2010, the highest coverage percentage was in the western region (Mouth of Puerto Real) with 20%, and in the southern part of the river discharge (zone 3) there are practically no seagrasses. The loss of vegetation cover in the southeast may be related to the low salinity recorded in the area (<10‰), as well as low water transparency (<20%). Since this vegetation is associated with high salinity and transparency values

and low levels of calcium carbonate (Yáñez-Arancibia et al., 1983; Raz-Guzman & Barba-Macías, 2000; Herrera Silveira et al., 2011 from Romo-Ríos, 2013)

Macroalgae

The distribution of macroalgae in Terminos Lagoon and the adjacent continental shelf is characterized by occurring in the sandy or sandy-silty substrate where algae such as *Cladophora vagabunda*, *Vauceheria sp.* and *Enteromorpha lingulata*. In the sediments of coarse grains, shells or shell fragments, the following species were recorded: *Gracilaria verrucosa*, *Spyridia filamentosa*, *Acanthophora spicifera*, *Hypnea cervicornis*, *H. cornuta*, *H. musciformis*, *Polysiphonia sp.*, *Enteromorpha lingulata*, *Aphanocapsa littoralis*; on large oyster shells: *Chondria baileyana*, *Gelidium pusillum* and *Dictyota ciliolata* (Conover, 1964; from Romo-Ríos, 2013).

Ortega (2009) mentions the importance of shells as a low specific weight algal substrate, which is why it constitutes a mobile community due to the action of waves and currents. The specifically algae that live on mangroves include *Bostrychia radicans*, *Caloglossa leprieurii* and *Microleus chthonoplastes*, although they are present in silts.

An important part of the vegetation is constituted by non-strict epiphytes, such as *Herposiphonia secunda*, *Polysiphonia ferulacea*, *Polysiphonia sp.*, *Heterosiphonia crispella*, *Centroceras calvulatum*, *Enteromorpha flexuosa* sub sp *paraxa*. Among the algae not fixed on some substrate or floating algae are *Caulerpa fastigiata* var. *confervoides*, *Chaetomorpha linum*, *Rhizoclonium africanum*, *R. kernerii* (Ortega, 2009).

Table II. Phytobenthic richness in Terminos Lagoon. (after Ortega, 1995).

Taxa		%	Genera	Species
Algae	Rhodophyceae	56.1	30	55
	Chlorophyceae	22.4	12	22
	Phaeophyceae	6.1	5	9
	Cyanophyceae	7.1	5	7
	Xanthophyceae	1	1	1
	Seagrass	Monocotyledoneae	5.1	4

Mangrove

Mangrove forests play an important role in the functioning of tropical coastal systems.

The four species reported for Mexico are present in Laguna de Terminos: *Rizophora mangle* or red mangrove in the margins of rivers, lagoons and coasts, usually flooded soils, *Laguncularia racemosa* or white mangrove found in flooded soils with high levels of salinity, *Avicennia germinans* or black mangrove on sandy soils that are flooded only

during part of the year or in years of maximum rainfall, *Conocarpus erecta* or buttonwood mostly on sandy and clay soils with low salinity that occasionally flooded during the rainy season (Agraz et al., 2012). In general, the forests that border the Terminos Lagoon present a 50% riparian appearance, 38.5% border, 10% basin and 1.5% scrub. This zoning is attributed to the heterogeneous environmental conditions that characterize the lagoon system, due to the fact that it receives large volumes of freshwater flows from the basin that drains into the Yucatán Peninsula, the lowlands of Tabasco and the Sierra de Chiapas. and Guatemala; as well as by the transport of terrigenic nutrients to the lagoon system, the type of soil and the anthropic activities that take place in the region of Terminos Lagoon (Agraz et al., 2012). It is relevant that the dominant appearance of the mangroves continues to be responsible for the high productivity and biodiversity that is currently recorded (Agraz et al., 2012). Various authors acknowledge that mangrove forests with riparian and border features are those that contribute the highest litter production, maintaining a rich and complex food chain characterized by high fish production (Agraz et al., 2012). On the other hand, the euryhaline water masses and the oxygen availability of the interstitial water from oxic to hypoxic prevailing in most of the Terminos Lagoon, determine the dominance of: black mangrove (*A. germinans*) with 77.4% presenting mechanisms of excretion, exclusion, accumulation of salt and the ability to thrive in conditions of low oxygen concentration. White mangrove (*L. racemosa*) presented a dominance of 22.2%, salt excretory glands; the red mangrove (*R. mangle*), is a species with low tolerance capacity to salinity and low oxygen concentrations (Agraz et al., 2012 in Romo-Ríos, 2013).

It has been observed that during the rainy and northern seasons, the largest volume of litter is contributed to the lagoon, which fertilizes the waters favoring increases in the biomass of consumers (Barreiro-Güemes, 1999; Agraz et al., 2012) .

In Terminos Lagoon, the main origin of organic matter is autochthonous with high contributions from the mangroves in the river discharge area both in the north and rainy season, so the degree of contribution of primary producers to Sedimentary Organic Matter in Terminos Lagoon, is directly related to the extension of the vegetation present in the system (Romo-Ríos, 2013)

The organic matter from the mangrove is an important component of the export of the lagoon system to the adjacent sea in the northern season due to the system's circulation pattern. (Romo-Ríos, 2013)

Fauna

This lagoon supports 117 species of fish, 69 of macro crustaceans (Raz, 2010), and 172 of mollusks (Reguero et al., 2010) among others. Of the species of fish and crustaceans

registered for the Terminos lagoon, 43 are commercially exploited (Ramos Miranda, 2000; Ramos Miranda et al., 2005a) (Ramos Miranda et al., 2015a)

Benthos

The Benthos of the lagoon reflects the circulation and sediment patterns. There is a pattern of distribution, diversity and frequency and it is possible to identify and characterize five faunal assemblages, highly correlated with salinity, substrate and primary producers in the different habitats: a) limnetic areas; b) fluvial-lagoon systems; c) inner lagoon; d) central basin and e) marine influenced areas. The species of commercial importance associated with these are: *Rangia* sp. in fluvial-lagoon sub-systems; *Crassostrea virginica* in inner lagoons (Pantanios); *Crassostrea rhizophora*, *Melongena melongena*, *Pleuroploca gigantea*, and *Strombus alatus* in marine influenced areas. Numerous marine and estuarine species of decapoda occur in Terminos Lagoon. The dominant species are those that can be termed slow-swimming like *Callinectes similis*, *C. sapidus*, and *C. bocourti* as well as four species of penaeid shrimp, *Penaeus setiferus* which is the most abundant in the southwestern area of the lagoon (fluvial-lagoon sub-system), *P. aztecus* in the west side near Carmen Inlet, *P. duorarum* in the east side of the lagoon near Puerto Real Inlet, and *Xiphopenaeus kroyeri* in the western side and in Carmen Inlet. The studies indicate that migration into the lagoon occurs through both inlets but mainly through Carmen Inlet. (Robadue, 2004)

After an analysis of the spatio-temporal behavior of benthos biodiversity the rainy season was the one that presented the highest values in terms of **diversity and richness indices**, while at the spatial level, region adjacent to the Mouth of Atasta and Palizada river (Region 1 after Villéger et al., 2010) was the most for both indices (Irola, et al., 2021). In terms of abundance, the internal coastline of Carmen Island (region 2 after Villéger et al., 2010) was the one characterized by presenting the highest values of abundance. Therefore, regions adjacent to the Mouth of Atasta and Palizada river and the internal coastline of Carmen Island (1 and 2) represent areas of great ecological importance for the balance of biodiversity, which is why they are key areas that should be protected in Terminos Lagoon. (Irola, et al., 2021)

Ichtyc community

The Terminos lagoon supports high biodiversity and together with the adjacent Campeche probe, along with the Campeche continental shelf maintains numerous commercially important fisheries which employ thousands of people in the southern Gulf of Mexico. It was recognized in 1994 as a Flora and Fauna Protection. It has been studied since the 1960s and historical data on fish communities and their abiotic conditions have been obtained since the 1970s.

Therefore, understanding the long-term dynamics of this resource is essential in determining the present community assemblage and current magnitude species

richness in order to support and maintain livelihoods in the region. (Paz-Ríos, et.al., 2021)

The spatial variations of the primary producers regulate the distribution of the fish and therefore the energy flow in terms of carbon and nitrogen of the organic matter in Terminos Lagoon and adjacent sea. The ichthyofauna present in the internal coastline of Carmen Island, Mouth of Puerto Real presents a trophic structure based on submerged vegetation (sea grasses and macroalgae) while the ichthyofauna of the fluvio-lagoon region (zone 3) presented a mangrove-based feeding source through detritus pathway. (Romo-Ríos, 2013)




Figure 7. Contribution of primary producers to ichthyofauna in “nortes”, after Romo-Ríos, 2013. (Z1, zone 1: Z2 zone 2, Z3, zone3, Z4, zone 4 PCA adyacent continental platform) and sample stations by Romo-Ríos, 2013




Figure 7. Contribution of primary producers to ichthyofauna in dry season, after Romo-Ríos, 2013. (Z1, zone 1: Z2 zone 2, Z3, zone3, Z4, zone 4 PCA adyacent continental platform)

Changes in the characteristics of the water have influenced the distribution of the species (Ramos-Miranda et al., 2015a); noting that the abundance of fish communities has decreased significantly over the years (Sosa-López et al., 2006; Ramos-Miranda et al., 2015; Flores-Hernández, et al., 2021), as well as a change in the dominance of

species (Ramos Miranda, et.al., 2015a), showing a greater change in its structure (biomass) than in the loss of species composition (Villegér, et al., 2010; Ramos Miranda, et.al., 2015a). The increase in salinity is probably one of the causes that can explain the variation in the abundance of the species. (Villegér, et al., 2010; Ramos Miranda, et.al., 2015a; Flores-Hernández, et al., 2021).

The behavior of the observed salinity, pointing out that the greater intrusion of seawater and the pattern of circulation in the lagoon ecosystem, have conditioned a reduction in the wide ranges of salinity observed in the 80's, and as a consequence an homogenization of the habitat (Flores-Hernández, 2021).

In the northern part of the lagoon, Villeger, et al., (2010) found an increase in fish richness but a significant decrease of functional divergence and functional specialization. They explained this result by a decline of specialized species (i.e., those with particular combinations of traits), while newly occurring species are redundant with those already present. The species that decreased in abundance have functional traits linked to seagrass habitats that regressed consecutively to increasing eutrophication (Villéger et al., 2010). Thus, Terminos Lagoon becomes a habitat more frequented by species of marine origin with a trend towards a more generalist fish community, under a regime more of sharing resources than of competition (Flores-Hernández, 2021).

Birds, reptiles and mammals

The most conspicuous fauna in the Terminos Lagoon are the Birds. The records for Terminos Lagoon (Programa de Manejo de Laguna de Terminos) indicate there are 49 families with 279 species in the complex. This area is considered to harbor at least during part of the year, 33% of the migratory birds moving along the Mississippi route using this area as feeding, sheltering and nesting habitat. Many of them are endangered and the most charismatic of these is the Jabiru Stork (*Jabiru micteria*). Recent studies indicate there are only 20 individuals left in Mexico. Other wetland birds and waterfowl like *Mycteria americana* (american stork), *Anas acuta*, *Anas cyanoptera*, *Mareca americana*, *Aythya affinis*, *Amazona albifrons* and *Chlorocere* sp among others. In terms of other fauna, the Reptiles found in this area are essentially associated with the freshwater wetlands of "Pantanos de Centla" on the western part of the Terminos Lagoon. Here significant populations of caiman (*Caiman crocodilus*) have been reported. This area holds areas where hawksbill turtle (*Eretmochelys imbricata*) and white turtle (*Chelonia mydas*) nest in the beaches of Terminos Lagoon. Because they are threatened species they hold special concern both for the Mexican authorities as well as world conservationists. Even though they are not easily seen, there are various Mammal species that use the Terminos Lagoon. Among them are dolphins (*Tursiops truncatus*) in the inlets connecting the lagoon with the Gula of Mexico oceanic waters. Other mammal species found here are more associated with wetland habitats

specifically the “Pantanos de Centla” are otters (*Lutra longicaudis*) and manatee (*Trichechus manatus*).-(Robadue, et al, 2014)

Socio-environmental characteristics

Terminos Lagoon has an important biodiversity, which has generated the idea of the possibility of intensively extracting only one of its resources until it is exhausted. In general with these productive and sociocultural practices, have prevailed in the extractive processes, demonstrating the lack of understanding of the importance of the coastal ecosystem as a reservoir of biocultural heritage. 12

In the region there are four economic, social and cultural processes that had been established linked to extensive extraction:

- a) Extraction of the tree “Palo de Campeche or Palo de Tinte” (*Haematoxylum campechianum*) for 250 years, which as a colorant had great importance in the European textile industry of the second half of the 18th century. until the 19th century, which implied struggles between Spain and England for control of the exploitation zones, among them Campeche, Laguna de Terminos and Tabasco (Contreras, 1990 in Villegas, et al., 2015);
- b) Later, 57 years of extensive and intensive exploitation of chicle, the chicle extraction industry generated a dynamic that marked the economy of the state of Campeche and the region of Terminos Lagoon during the first forty years of the twentieth century, although this activity has been recognized since 1890 (Vadillo, 2001 in Villegas, et al., 2015).

Both processes allowed the reshaping of the regional oligarchy closely linked to the US commercial capital, contributing during its exploitation to the deforestation of the Campeche forests (Vadillo, 2001 in Villegas, et al., 2015);

- c) Subsequently, in 50 years of intensive shrimp exploitation in El Carmen, its origins, evolution and situation at the time of the transfer of the shrimp fleet from the private sector to the cooperatives in 1982, represents an outward-oriented development model, although their degree of vulnerability and dependency are lower by virtue of the provisions of the Federal Government that stimulated the growth of an industry and promoted in the locality the establishment of manufacturing institutions to process, freeze and pack shrimp, and to build and repair fishing vessels (Leriche, 2001 in Villegas, et al., 2015);
- d) And until now a days, the last extractive stage with 35 years of intensive exploitation of hydrocarbons and gas with the installation of platforms in the marine area. These two social events during the twentieth century: the arrival of deep-sea fishing and later the oil activity, both are confronted with the millenary riverine culture of the region, when presenting themselves as

modernization processes give rise to conflicts and tensions between the different social groups –those who inhabit and those who arrive– that are confronted by the appropriation of the territory and with it in time their spatial and sociocultural reconfiguration is observed (Villegas, 2008 in Villegas, et al., 2015).

Blue crab habitat

The life history of the blue crab involves a complex cycle of planktonic, nektonic, and benthic stages which occur throughout the estuarine-nearshore marine environment, occupying a variety of habitats depending on the requirements of each life history stage (Perry et al., 1984).

Its preferred habitat is the shallow coastal areas of the estuarine complex, with muddy and mud-clay bottoms, but they support a wide variation in environmental conditions, from fresh water (0 ‰) to hypersaline lagoons, temperature ranges ranging from 3 °C up to 35 °C and are also present in shallow waters or at depths of 90 m.
[\(https://www.ecured.cu/Jaiba_azul\)](https://www.ecured.cu/Jaiba_azul)

Female blue crabs are catadromous; they migrate from hyposaline waters to higher salinity water to spawn and hatch their eggs. They are commonly seen in September-October offshore Terminos Lagoon (PescadoresXXX)- The high salinity, oceanic water not only serves as habitat for the spawning female but ensures larval development, increases dispersal capabilities, decreases osmoregulatory stress, and reduces predation. Fertile eggs hatch into free swimming larvae (zoeae) which pass through a series of molts. Newly-hatched blue crab larvae normally develop through seven zoeal stages before transforming into a megalopal stage. (VanderKooy, 2013)

Megalopae return to the estuary where they metamorphose into the first crab stage. The estuarine phase is perhaps the most critical because all post-settlement growth and the major components of the reproductive cycle occur there. Male blue crabs usually remain within the estuary during their entire post-settlement life. Juvenile and adult blue crabs exhibit wide seasonal and areal distribution within estuaries.

Among the factors that limit its growth and survival are the low availability of food, the predation of other organisms, the deterioration of the habitat, since it limits its shelter and feeding and the accumulation of pollutants in the sediment given its benthic nature, while that within the physical and chemical environmental parameters of water, the one that has the greatest influence on its distribution is salinity, high temperatures and the decrease in dissolved oxygen. (VanderKooy, 2013).

Other environmental parameters that influence its abundance and distribution are rainfall and droughts due to their direct relationship with salinity and strong meteorological events due to habitat destruction.

Monitoring Program FIP Results

50 interviews were carried out with fishermen and suppliers who deliver to the FIP during 2023, using the record attached at the end of this report. When possible interviews were made as observing the by-catch on the boats. Throughout the work of the FIP we have sought to ensure that the fishermen who collaborate with us understand the need and usefulness of taking care of the habitat and the ecosystem, keeping their spaces free of garbage, recycling, releasing the organisms in the fishing ground alive and in good conditions, and promoting to comply with established regulations. Over the years, the percentage of blue crabs below the minimum legal size has remained less than 5% in total. The observed and reported bycatch have decreased, as fishers release alive organisms in the fishing ground. Catches of *Menippe mercenaria* are sometimes present (chelaes), but in lesser extent. An increase in the number of soft crab buyers has been observed in the localities. The FIP, not being able to compete with the income it generates for fishermen, is promoting the release of at least the juvenile females.

Palo Alto

Fishermen in this area, most of the year, fish in canoes propelled by wooden poles, without a motor. Their catches per day vary between 18 and 35 kilos of crab per day. They make their catches with cages that they leave working in the evening and collect in the early morning of the next day. By not requiring to pay for gasoline, its captures are economically efficient. Even so, buyers of smooth crab have been detected in their landing area. They catch their catches in shallow areas of between half a meter and one meter in muddy, grassy or sandy areas. They commonly do not bring associated catches on their boats except for the fish checkered puffer (*Sphoeroides testudineus*) and some crabs that they release in the mangrove and species for self-consumption.

Los pescadores de esta zona, la mayor parte del año pescan en cayucos impulsados por palos de madera, sin motor. Sus capturas por día varían entre 18 y 35 kilos de jaiba por día. Realizan sus capturas con jaulas que dejan trabajando por la tarde-noche y recogen en la madrugada del día siguiente. Al no requerir pagar gasolina, sus capturas son económicamente eficientes. Aun así, se ha detectado a compradores de jaiba suave en su zona de desembarco. Sus capturas las realizan en zonas poco profundas de entre medio metro y un metro en zonas lodosas, con pastos o arenosas. Comúnmente no traen en sus embarcaciones capturas asociadas salvo the fish checkered puffer

(*Sphoeroides testudineus*) y algunos cangrejos que liberan en el manglar y especies para autoconsumo.

Isla Aguada

Purchases are made through a permit holder, so it is not possible to verify the species they bring on their boats. Periodically they are given short talks about the importance of taking care of the habitat of the species they capture. These fishermen go out with motor boats, generally catching between half a meter and 3 meters deep, in seagrass or muddy-sandy areas. Since they recognize the importance of the habitat, and that they use to fish in the same area for several years (some of them for around 10 years), they usually leave non commercial or bait species in the fishing ground. When there is bad climate conditions they sometimes bring by catch back to the coast.

Las compras se realizan a través de una permisionaria, por lo cual no se puede constatar las especies que traen en sus embarcaciones. Periódicamente se les da pequeñas charlas sobre la importancia de cuidar el hábitat de las especies que capturan. Estos pescadores salen con embarcaciones con motor, generalmente capturan entre medio metro y 3 metros de profundidad, en zonas de pastos marinos o lodoso-arenosas.

Atasta

In this case, purchases are also made from a permit holder. Depending on the time of year, fishermen go fishing in cayuco (during the rainy season) propelled by poles, capturing in flood-prone areas. Catches in this area are made at depths of less than one meter, and generally between 30 and 50 cm. *C. rathbunae* is frequently caught in these areas. Fishermen do not usually catch other species. In the dry season they go fishing in motor boats at depths between half a meter and 3 meters. Most of these catches are of the species *C. sapidus*, generally reporting mojarras, catfish, and seabream in their catches. Some crustaceans are released alive either in the fishing area or on the banks of the mangrove. Most of these fishermen look for seagrass beds to fish, although they comment that these have been disappearing, so they also tend to fish in muddy beds. In order to save money, some of them go out to fish with nets, which will be their bait.

En este caso las compras también se realizan a partir de una permisionaria. Dependiendo de la época del año los pescadores salen a pescar en cayuco (durante la época de lluvias) impulsadas por palos capturando en zonas inundables. Las capturas de esta zona se realizan a profundidades menores a un metro, y generalmente entre 30 y 50 cms. En estas zonas con frecuencia se captura *C. rathbunae*. Los pescadores no

suelen capturar otras especies. En época de secas salen a pescar en embarcaciones con motor a profundidades entre medio metro y 3 metros. La mayor parte de estas capturas son de la especie *C. sapidus*, generalmente reportan en sus capturas mojarras, catfish, and seabream. Algunos crustáceos son liberados con vida ya sea en la zona de pesca o en las orillas del manglar. La mayoría de estos pescadores buscan los fondos de pastos marinos para pescar aunque comentan que estos han ido desapareciendo, así que también suelen pescar en fondos lodosos.

Sabancuy

These fishermen deliver directly to the receiving area of the processing plant. They generally fish in very shallow areas when they cannot access other types of fishing resources. You cannot observe the accompanying fauna although they mention that they prefer to release everything that is not useful to them in the fishing site to ensure that there are crabs. They have maintained their fishing sites for many years and mention that this practice has allowed them to continue fishing.

Estos pescadores entregan directamente en el área de recepción de la planta procesadora. Generalmente pescan en áreas muy poco profundas cuando no pueden acceder a otro tipo de recursos pesqueros. No se puede observar la fauna acompañante aunque ellos mencionan que prefieren liberar todo lo que no les sirve en el sitio de pesca para asegurar que haya jaiba. Sus sitios de pesca los han mantenido por muchos años y mencionan que esa práctica les ha permitido seguir pescando.

Evidence of no or limited bycatch in the boats



Baskets without bycatch



Baskets without bycatch



Bait purchase



Baskets without bycatch





Literature cited

- Agraz, H., C.M., Osti S, J., Chan K, C.A., Gómez R, D., Castillo D, S., Expósito D, G., Márquez, É., Acosta, J., Reyes C, J. & Conde M, K.P. 2012. Diagnóstico para la Conservación y Restauración de esta Área de Protección de Flora y Fauna
- Barreiro-Güemes, M.T. 1999. Aporte de hojarasca y renovación foliar del manglar en un sistema estuarino del Sureste de México. Revista de Biología Tropical, 47: 729–737.
- Borges Souza, G.N. 2004. Composición y distribución de sedimentos en la Laguna de Términos, Campeche, México (2001-2002). Tesis de licenciatura. Universidad Autónoma de Campeche Facultad de Ciencias Químico-Biológicas.
- Conover, J.T. 1964. The Ecology, Seasonal Periodicity, and Distribution of Benthic Plants in some Texas Lagoons. Botanica Marina, 7: 4–41.
- Contreras, A., 1990. Historia de una tintórea olvidada. El proceso de explotación y circulación del palo de tinte 1750 – 1807. México: Universidad Autónoma de Yucatán.
- David, L. T., and B. Kjerfve. 1998. Tides and currents in a two inlet coastal lagoon: Laguna de Terminos, Mexico. Continental Shelf Research 18:1057–1079
- Escobar-Toledo F., Zetina-Rejón M. J., Ramos-Miranda J., Arreguín-Sánchez F. Temporal shifts in functional traits of the fish community in Terminos Lagoon (Mexico) in three periods (1980, 1998 and 2011. Environmental Biology of Fishes. 2017;100:1575–1586. doi: 10.1007/s10641-017-0666-2.
- Feria Alvarado, K.G., R.F. Chávez González, Y.E. Torres Rojas y R.E. del Río Rodríguez, 2021. Morphological aspects and its influence in the ecological role of two sympatric species (*Lagodon rhomboides* and *Archosargus rhomboidalis*) in Terminos lagoon, Mexico. JAINA Costas y Mares ante el Cambio Climático, 3(1): 41-54. doi 10.26359/52462.0321
- Flores Hernández, D., J. Ramos Miranda, J. Rendón von Osten, L.A. Ayala Pérez, A. Sosa López, L. Alpuche Gual, R. Rosa Vega, F. Arreguín Sánchez, M.E. Gonzalez y de la Rosa, y J. Santos Valencia, 2001. Evaluación del camarón blanco (*Litopenaeus setiferus*) y de las comunidades neóticas de la Laguna de Términos, Campeche, México: estimación de los impactos ambientales y pesqueros. alim-11- 96. sisierra-uac-anplt. 19 p.
- Flores-Hernández, D., J. Ramos-Miranda, L. A. Ayala-Pérez y F. Gómez Criollo, 2021. Environmental variation vs co-occurrence of two fish families (Gerridae and Ariidae) in Terminos Lagoon, Campeche, Mexico. JAINA Costas y Mares ante el Cambio Climático, 3(1): 5-22. doi 10.26359/52462.0121
- Gold-Bouchot, G., Barroso-Noreña, E., Zapata-Pérez, O., 1995. Hydrocarbon concentrations in the american oyster (*Crassostrea virginica*) in Laguna de Terminos,

Campeche, Mexico. Bulletin of Environmental Contamination and Toxicology, 53(2): 222-227.

Herrera Silveria, J., Silva Casarín, A., Salles Alfonso De Almeida, P., Villalobos Zapata, G.J., Medina Gomez, I., Espinal González, J.C., Zaldívar Jiménez, A., Trejo Peña, J., González Jáuregui, M. & Cu Escamilla, A.D. 2002. Análisis de Calidad Ambiental usando indicadores hidrobiológicos y modelo hidrodinámico actualizado de la Laguna de Términos (CINVESTAV-IPN U. Mérida, EPOMEX-UAC; Inst. Ingeniería -UNAM).

Herrera-Silveira, J.A., Arellano Méndez, L., & Morales Ojeda, S. 2011. Report of the Pilot Project Natural Habitat and Ecosystem Conservation of Coastal and Marine Zones of the Gulf of Mexico: Seagrass beds characterizatio

Irola-Sansores, B.N., Y.E. Torres Rojas, y A. Cuevas Jiménez, 2021. Spatial-temporal behavior of benthic communities in the Terminos Lagoon, Mexico: Possible areas of protection and conservation. JAINA Costas y Mares ante el Cambio Climático, 3(1): 55-76. doi 10.26359/52462.0421

Leriche, L., 2001. Isla del Carmen: La historia indecisa de un puerto exportado. El caso de la industria camaronera (1947-1982). México: Gobierno del Estado de Campeche. Universidad Autónoma del Carmen, Instituto de Cultura de Campeche.

Moore, K.A. & L. Wetzel, Y.R. 1988. The distribution and productivity of seagrasses in the Términos Lagoon. In: Ecología de los ecosistemas costeros en el sur del Golfo de México: La región de la Laguna de Términos A. Yáñez Arancibia & J.W. Day Jr. Inst. Cienc. Del Mar y Limnol. UNAM. Coast. Ecol. Inst. LSU. Editorial Universitaria México DF Cap. 12:207-220.

Ortega, M.M. 2009. Observaciones del fitobentos de la Laguna de Términos, Campeche, México. Anales del Instituto de Biología serie Botánica, 66.

Paz-Ríos, C. E., Atahualpa Sosa-López, corresponding author¹ Yassir E. Torres-Rojas,¹ Julia Ramos-Miranda,¹ and Rodolfo E. del Río-Rodríguez¹ Fish species richness in the Terminos Lagoon: An occurrence data compilation of four sampling campaigns along a multidecadal series Biodivers Data J. 2021; 9: e65317 Published online 2021 May 21. doi: 10.3897/BDJ.9.e65317

Ramos Miranda J., y G.J. Villalobos Zapata (editores), 2015. Aspectos socioambientales de la región de la laguna de Términos, Campeche. Universidad Autónoma de Campeche. 210 p.

Ramos Miranda, J., Flores Hernández, D., Sosa López, A., Ayala Pérez, L., Villéger, S., Pérez Sánchez, M. I., Gómez Criollo, F., Can González, M., Flores Ramos, E. F. y Mouillot, D. 2015a. La comunidad del necton en la laguna de Términos: modificación del hábitat vs cambios en la comunidad. Universidad Autónoma de Campeche.

Ramos Miranda, J., L. Quiniou, D. Flores Hernández, T. Do Chi, L.A. Ayala Pérez, & A. Sosa López, 2005b. Spatial and temporal changes in the nekton communities of Terminos Lagoon Campeche, Mexico. *Journal of Fish Biology*, 66, 513- 530.

Ramos Miranda, J., Mouillot, D., Sosa Lopez, A., Do Chi, T. & Flores Hernández, D. 2008. How much variation can be explained by seasonal, spatial and environmental effects in nekton assemblages of the Terminos Lagoon? *Aquatic Conservation: Marine and Freshwater Ecosystems*, 18: 508–517.

Ramos-Miranda J., Mouillot D., Flores-Hernández D., Sosa-López A., Do-Chi T., Ayala-Pérez L. A. 2005a Changes in four complementary facets of fish diversity in a tropical coastal lagoon after 18 years: a functional interpretation. *Marine Ecology Progress Series*. 2005a ;304:1-13. doi: 10.3354/meps304001

Raz-Guzmán, A. & Barba-Macias, E. 2000. Seagrass biomas, distribution and associated macrofauna in southwestern of Mexico coastal lagoons. *Biol. Mar. Mediterranean*, 7(2): 271.

Rendón-von-Osten, J., Memije-Canepa, M., Ek-Moo, N.A., 2005. Plaguicidas orgánicos persistentes (POPs) en sedimentos de la costa sur de Campeche, México. p. 249- 260. In: Botello AV, Rendón von Osten J, Bouchot G y Agraz Hernández C (eds.). Golfo de México Contaminación e Impacto Ambiental: Diagnóstico y Tendencias, 2da. Edición. Univ. Autón. De Campeche, Univ. Nal. Autón. de México, Instituto Nacional de Ecología.

Rivera-Arriaga E., Villalobos-Zapata G. J. The coastal zone of Campeche, Mexico: Opportunities for implementing and integrated coastal management framework. *Journal of Coastal Research*. 2005;42:184–190. [Google Scholar]

Robadue, D., Oczkowski, A., Calderon, R., Bach, L., & Cepeda, M. F. 2004. Characterization of the Region of the Términos Lagoon: Campeche, Fexico: Draft for discussion. Narragansett, RI; Coastal Resources Center, University of Rhode Island. PLUS Calderon, R. (2004) Draft 1 Site Profile: The Laguna de Términos, México. Corpus Christi, TX: The Nature Conservancy.

Romo-Ríos, J.A. 2013. Productores primarios y su relación con la distribución de la ictiofauna en Laguna de Términos y Plataforma Continental adyacente, Campeche, México. Tesis para obtener el grado de Maestría en Ciencias en Manejo de Recursos Marinos. CICIMAR-IPN. 84pp.

Santos Santoyo, J.E., Y. E. Torres Rojas, y A. Cuevas Jiménez, 2021. Record of changes in the regionalization of Terminos Lagoon, Campeche. *JAINA Costas y Mares ante el Cambio Climático*, 3(1): 23-40. doi 10.26359/52462.0221

SEMARNAP-INE 1997. Programa de Manejo del Área de Protección de Flora y Fauna Laguna de Términos, México. SEMARNAP, México D.F., 166

Sosa López A., D. Mouillot, T. Do Chi, & J. Ramos Miranda, 2005. Ecological indicators based on fish biomass distribution along trophic levels: an application to the Terminos coastal lagoon, Mexico. Ices Journal of Marine Science, 62: 453-458.

Sosa López, A., D. Mouillot, J. Ramos Miranda, D. Flores Hernández & T. Do Chí, 2007. Fish richness decreases with salinity in tropical coastal lagoons Journal of Biography, 34: 52-6

Vadillo, C., 2001. Los chicleros en la región de Laguna de Términos, Campeche: 1890 – 1947. México: Universidad Autónoma del Carmen

Villegas , S. J. y Adriana, Solís F. 2015. Crisis ambiental, áreas naturales progenidas y turismo. Facultad de Ciencias Sociales Económico Administrativas, Universidad Autónoma del Carmen. 41-64 pp. en Ramos Miranda J., y G.J. Villalobos Zapata (editores), 2015. Aspectos socioambientales de la región de la laguna de Términos, Campeche. Universidad Autónoma de Campeche. 210 p.

Villegas, J., 2008. Organización social, modernización y utopías entre los pescadores ribereños de la Península de Atasta e Isla Aguada, Campeche. En: Fraga, J.; Villalobos, G.; Doyon, S. y García, A. (Eds.) Descentralización y manejo ambiental. Gobernanza costera en México. idrc, crdi, Cinvestav, Epomex, Plaza y Valdés.

Villéger, S., J. Ramos Miranda, D. Flores Hernández, & y D. Mouillot, 2010. Contrasted changes in taxonomic versus functional diversity of tropical fish communities after habitat degradation. Ecological Applications, 20(6): 1512-1522.

Yañez-Arancibia et al. 1982 defined environmental zones after environmental condiciones recorded in 1980-1981 including depth, water transparency, bottom salinity and substrate characteristics defining 4 zones. . Another similar biological survey were conducted in 1998-1999 (Ramos Miranda, 2000).

"Yáñez-Arancibia, A. & J. W. Day, Jr. 1982. Ecological characterization of Términos Lagoon, a tropical lagoon-estuarine system in the Southern Gulf of Mexico, p. 431-440. In: Laserre, P. y H. Postma (Eds.) Costal Lagoons. Oceanologica Acta Vol. Spec., 5 (4): 462 p"

Yáñez-Arancibia, A., Lara-Domínguez, A., Chavance, P. & Flores Hernández, D. 1983. Environmental Behavior of Terminos Lagoon Ecological System, Campeche México. An. Inst. Cien. Del Mar Y Limnol. UNAM, 10 (1): 137–176.

Annex 1. Systematic list of species captured in Laguna de Terminos, period 2016-2017 (taken from Irola, et al., 2021)

	Family	Species
Bivalva	Ostreidae	<i>Crassostrea virginica</i>
Gastrópoda	Cassidae	<i>Semicassis granulata</i>
	Fasciolariidae	<i>Triplofusus giganteus</i>
	Buccinidae	<i>Sinistrofulgur perversum</i>
		<i>Fulguropsis spirata</i>
	Strombidae	<i>Strombus alatus</i>
Cephalopoda	Loliginidae	<i>Lolliguncula brevis</i>
	Squillidae	<i>Squilla empusa</i>
Crustacea	Peneidae	<i>Penaeus aztecus</i>
		<i>Penaeus (Farfantepenaeus) duorarum</i>
		<i>Penaeus (Litopenaeus) setiferus</i>
		<i>Rimapenaeus similis</i>
		<i>Xiphopenaeus kroyeri</i>
	Sicyoniidae	<i>Sicyonia brevirostris</i>
	Menippidae	<i>Menippe mercenaria</i>
	Portunidae	<i>Callinectes bocourti</i>
		<i>Callinectes danae</i>
		<i>Callinectes rathbunae</i>
		<i>Callinectes sapidus</i>
		<i>Callinectes similis</i>
Rayas	Rhinobatidae	<i>Pseudobatos lentiginosus</i>
	Dasyatidae	<i>Dasyatis sabina</i>
		<i>Himantura schmardae</i>
		<i>Hypanus americanus</i>
	Urotrygonidae	<i>Urobatis jamaicensis</i>
Teleosti	Gymnuridae	<i>Gymnura micrura</i>
	Clupeidae	<i>Dorosoma analis</i>
		<i>Harengula jaguana</i>
		<i>Opisthonema oglinum</i>
	Engraulidae	<i>Anchoa hepsetus</i>
		<i>Anchoa lamprotaenia</i>
		<i>Anchoa mitchilli</i>
		<i>Cetengraulis edentulus</i>
	Synodontidae	<i>Synodus foetens</i>
	Ariidae	<i>Ariopsis felis</i>
		<i>Arius melanopus</i>

	<i>Bagre marinus</i>
Batrachoididae	<i>Opsanus beta</i>
Hemiramphidae	<i>Chriodus atherinoides</i>
	<i>Hippocampus zosterae</i>
	<i>Hippocampus erectus</i>
Syngnathidae	<i>Syngnathus floridae</i>
	<i>Syngnathus louisianae</i>
	<i>Syngnathus scovelli</i>
	<i>Prionotus carolinus</i>
Triglidae	<i>Prionotus scitulus</i>
	<i>Prionotus tribulus</i>
Serranidae	<i>Diplectrum formosum</i>
	<i>Caranx hippos</i>
Carangidae	<i>Chloroscombrus chrysurus</i>
	<i>Hemicaranx amblyrhinchus</i>
	<i>Selene vomer</i>
	<i>Lutjanus analis</i>
Lutjanidae	<i>Lutjanus griseus</i>
	<i>Lutjanus synagris</i>
	<i>Diapterus auratus</i>
Gerreidae	<i>Diapterus rhombeus</i>
	<i>Eucinostomus argenteus</i>
	<i>Eucinostomus gula</i>
	<i>Eucinostomus melanopterus</i>
	<i>Eugerres plumieri</i>
	<i>Stenotomus caprinus</i>
	<i>Anisotremus virginicus</i>
Pomadasytidae	<i>Conodon nobilis</i>
	<i>Haemulon aurolineatum</i>
	<i>Haemulon bonariense</i>
	<i>Haemulon plumieri</i>
	<i>Orthopristis chrysoptera</i>
	<i>Archosargus probatocephalus</i>
Sparidae	<i>Archosargus rhomboidalis</i>
	<i>Lagodon rhomboides</i>
	<i>Bairdiella chrysura</i>
Sciaenidae	<i>Bairdiella ronchus</i>
	<i>Corvula batabana</i>
	<i>Cynoscion arenarius</i>
	<i>Cynoscion nebulosus</i>

	<i>Cynoscion nothus</i>
	<i>Menticirrhus americanus</i>
	<i>Menticirrhus saxtilis</i>
	<i>Menticirrhus littoralis</i>
	<i>Micropogonias undulatus</i>
	<i>Stellifer lanceolatus</i>
Polynemidae	<i>Polydactylus octonemus</i>
Cichlidae	<i>Cichlasoma urophthalmus</i>
Gobiidae	<i>Gobioides broussonnetti</i>
	<i>Gobionellus oceanicus</i>
Ephippidae	<i>Chaetodipterus faber</i>
Trichiuridae	<i>Trichiurus lepturus</i>
Bothidae	<i>Citharichthys spilopterus</i>
Soleidae	<i>Achirus lineatus</i>
	<i>Trinectes maculatus</i>
Cynoglossidae	<i>Sympodus civitatum</i>
	<i>Sympodus plagiusa</i>
Monacanthidae	<i>Aluterus schoepfii</i>
	<i>Monacanthus ciliatus</i>
Ostraciidae	<i>Stephanolepis hispidus</i>
Tetraodontidae	<i>Acanthostracion quadricornis</i>
	<i>Sphoeroides greeleyi</i>
	<i>Sphoeroides nephelus</i>
	<i>Sphoeroides parvus</i>
	<i>Sphoeroides spengleri</i>
	<i>Sphoeroides testudineus</i>
Diodontidae	<i>Chilomycterus schoepfi</i>

Annex 2. Fish species richness in the Terminos Lagoon: An occurrence data compilation of four sampling campaigns along a multidecadal series. From Paz-Ríos, et al., 2021.

Class	Order	Family	Species	1980-1981	1998-1999	2010-2011	2016-2017
Elasmobranchii	Myliobatiformes	Dasyatidae	<i>Dasyatis hastata</i> (DeKay, 1842)	•			
			* <i>Hynpanus americanus</i> (Hildebrand & Schroeder, 1928)			•	•
			~ <i>Hynpanus sabinus</i> (Lesueur, 1824)	•	•	•	•
		Gymnuridae	<i>Gymnura micrura</i> (Bloch & Schneider, 1801)			•	•
		Potamotrygonidae	<i>Styracura schmardae</i> (Werner, 1904)		•		•
	Rhinopristiformes	Urotrygonidae	<i>Urobatis jamaicensis</i> (Cuvier, 1816)	•	•	•	•
		Rhinobatidae	<i>Pseudobatos lentiginosus</i> (Garman, 1880)			•	•
		Ephippidae	*~ <i>Chaetodipterus faber</i> (Broussonet, 1782)	•	•	•	•
		Lobotidae	*~ <i>Lobotes surinamensis</i> (Bloch, 1790)		•		
		Muraenidae	<i>Gymnothorax nigromarginatus</i> (Girard, 1858)			•	
Actinopteri	Anguilliformes		<i>Gymnothorax saxicola</i> Jordan & Davis, 1891		•		
		Ophichthidae	<i>Ophichthus gomesii</i> (Castelnau, 1855)	•	•		
	Aulopiformes	Synodontidae	<i>Synodus foetens</i> (Linnaeus, 1766)	•	•	•	•
		Batrachoidiformes	*~ <i>Opsanus beta</i> (Goode & Bean, 1880)	•	•	•	•
		Batrachoididae	<i>Porichthys porosissimus</i> (Cuvier, 1829)	•	•	•	
	Beloniformes	Belonidae	~ <i>Strongylura notata</i> (Poey, 1860)		•		
		Hemiramphidae	~ <i>Chriodorus atherinoides</i> Goode & Bean, 1882	•			•
		Blenniiformes	<i>Hyleurochilus geminatus</i> (Wood, 1825)				•
		Labrisomidae	<i>Paraclinus fasciatus</i> (Steindachner, 1876)				•
		Acaridae	~ <i>Achirus lineatus</i> (Linnaeus, 1758)	•	•	•	•
	Carangiformes		*~ <i>Trinectes maculatus</i> (Bloch & Schneider, 1801)	•	•	•	•
		Bothidae	<i>Bothus ocellatus</i> (Agassiz, 1831)	•			
		Carangidae	*~ <i>Caranx cryos</i> (Mitchill, 1815)		•	•	
			*~ <i>Caranx hippos</i> (Linnaeus, 1766)	•	•	•	•
			~ <i>Caranx latus</i> Agassiz, 1831			•	
			<i>Caranx ruber</i> (Bloch, 1793)			•	
			*~ <i>Chloroscombrus chrysurus</i> (Linnaeus, 1766)	•	•	•	•
			<i>Hemicarax amblyrhynchus</i> (Cuvier, 1833)		•	•	•
			~ <i>Oligoplites saurus</i> (Bloch & Schneider, 1801)	•	•	•	
			*~ <i>Selene setapinnis</i> (Mitchill, 1815)		•	•	
			<i>Selene vomer</i> (Linnaeus, 1758)	•	•	•	•
			*~ <i>Trachinotus carolinus</i> (Linnaeus, 1766)		•		
			<i>Trachinotus falcatus</i> (Linnaeus, 1758)		•		
			<i>Trachinotus goodei</i> Jordan & Evermann, 1896		•		
Clupeiformes	Centropomidae		*~ <i>Centropomus parallelus</i> Poey, 1860		•	•	
			*~ <i>Centropomus poeyi</i> Chávez, 1961		•		
			*~ <i>Centropomus undecimalis</i> (Bloch, 1792)	•			
	Cynoglossidae		*~ <i>Syphurus civitatum</i> Ginsburg, 1951		•		•
			*~ <i>Syphurus plagiusa</i> (Linnaeus, 1766)	•	•	•	•
	Paralichthyidae	Anclopussetta quadrocellata	Gill, 1864	•		•	
			~ <i>Citharichthys spilopterus</i> Günther, 1862	•	•	•	•
			<i>Etropus crossotus</i> Jordan & Gilbert, 1882	•	•	•	•
			<i>Syacium gunteri</i> Ginsburg, 1933			•	
	Polynemidae	<i>Polydactylus octonemus</i> (Girard, 1858)		•	•	•	•
		<i>Sphyraena barracuda</i> (Edwards, 1771)				•	
	Cichliformes	Cichlidae	~ <i>Mayaheros urophthalmus</i> (Günther, 1862)	•	•	•	•
		Clupeidae	*~ <i>Brevoortia gunteri</i> Hildebrand, 1948		•		
	Engraulidae		*~ <i>Dorosoma analle</i> Meek, 1904			•	
			~ <i>Dorosoma petenense</i> (Günther, 1867)		•	•	
			*~ <i>Harengula clupeola</i> (Cuvier, 1829)				•
			*~ <i>Harengula jaguana</i> Poey, 1865	•	•	•	•
			*~ <i>Opisthonema oglinum</i> (Lesueur, 1818)	•	•	•	•
			*~ <i>Sardinella aurita</i> Valenciennes, 1847		•		
			~ <i>Anchoa hepsetus</i> (Linnaeus, 1758)	•	•	•	•
			<i>Anchoa lamprotaenia</i> Hildebrand, 1943	•			•
			<i>Anchoa lyolepis</i> (Evermann & Marsh, 1900)			•	
			~ <i>Anchoa mitchilli</i> (Valenciennes, 1848)	•	•	•	•
			<i>Cetengraulis edentulus</i> (Cuvier, 1829)	•	•	•	•

Class	Order	Family	Species	1980-1981	1998-1999	2010-2011	2016-2017
<i>Actinopteri</i>	Cyprinodontiforme	Fundulidae	~Lucania parva (Baird & Girard, 1855)			●	
	Elopiformes	Elopidae	Elops saurus Linnaeus, 1766		●		
	Gobiiformes	Eleotridae	~Eleotris pisonis (Gmelin, 1789)				●
		Gobiidae	~Bathygobius soporator (Valenciennes, 1837)		●		
			~Gobionellus oceanicus (Pallas, 1770)	●	●	●	●
			Gobiosoma longipala Ginsburg, 1933				●
	Mugiliformes	Mugilidae	*~Mugil cephalus Linnaeus, 1758		●		
			*~Mugil curema Valenciennes, 1836		●	●	
	Perciformes	Scaridae	Nicholsina usta (Valenciennes, 1840)	●	●		
		Scorpaenidae	Scorpaena brasiliensis Cuvier, 1829			●	
			Scorpaena plumieri Bloch, 1789	●		●	●
		Triglidae	Prionotus beanii Goode, 1896	●			
			Prionotus carolinus (Linnaeus, 1771)	●	●	●	
			Prionotus martis Ginsburg, 1950			●	
			Prionotus punctatus (Bloch, 1793)	●		●	
			Prionotus rubio Jordan, 1886			●	●
			Prionotus scitulus Jordan & Gilbert, 1882	●	●	●	●
			Prionotus tribulus Cuvier, 1829	●		●	●
		Gerreidae	~Diapterus auratus Ranzani, 1842		●	●	●
			~Diapterus rhombeus (Cuvier, 1829)	●	●	●	●
			~Eucinostomus argenteus Baird & Girard, 1855	●	●	●	●
			~Eucinostomus gula (Quoy & Gaimard, 1824)	●	●	●	●
			~Eucinostomus melanopterus (Bleeker, 1863)	●	●		●
			*~Eugerres plumieri (Cuvier, 1830)	●	●	●	●
		Haemulidae	Anisotremus virginicus (Linnaeus, 1758)				●
			*~Conodon nobilis (Linnaeus, 1758)		●		
			Haemulon aurolineatum Cuvier, 1830		●	●	●
			*Haemulon bonariense Cuvier, 1830	●	●	●	●
			*Haemulon plumieri (Lacepède, 1801)	●	●	●	●
			*~Orthopristis chrysoptera (Linnaeus, 1766)	●	●	●	●
		Lutjanidae	*~Lutjanus analis (Cuvier, 1828)	●	●		●
			*Lutjanus apodus (Walbaum, 1792)		●		
			*~Lutjanus griseus (Linnaeus, 1758)	●	●	●	●
			*Lutjanus synagris (Linnaeus, 1758)	●	●	●	●
			*Ocyurus chrysurus (Bloch, 1791)		●		
		Sciaenidae	*~Bairdiella chrysoura (Lacepède, 1802)	●	●	●	●
			*Bairdiella ronchus (Cuvier, 1830)	●	●	●	●
			Corvula batabana (Poey, 1860)				●
			*~Cynoscion arenarius Ginsburg, 1930	●	●	●	●
			*~Cynoscion nebulosus (Cuvier, 1830)	●	●	●	●
			*Cynoscion nothus (Holbrook, 1848)	●		●	●
			*Menticirrhus americanus (Linnaeus, 1758)	●	●	●	●
			*Menticirrhus littoralis (Holbrook, 1847)				●
			Menticirrhus saxatilis (Bloch & Schneider, 1801)	●	●	●	●
			*Micropogonias furnieri (Desmarest, 1823)			●	●
			*~Micropogonias undulatus (Linnaeus, 1766)	●	●	●	●
			Odontoscion dentex (Cuvier, 1830)	●			
		Serranidae	*~Stellifer lanceolatus (Holbrook, 1855)	●	●	●	●
			Diplectrum bivittatum (Valenciennes, 1828)			●	
			*Diplectrum formosum (Linnaeus, 1766)			●	●
			Epinephelus itajara (Lichtenstein, 1822)	●			
		Sparidae	*Mycteroperca bonaci (Poey, 1860)		●	●	
			*~Archosargus probatocephalus (Walbaum, 1792)	●	●	●	●
			Archosargus rhombooidalis (Linnaeus, 1758)	●	●	●	●
			Calamus penna (Valenciennes, 1830)		●		
			*~Lagodon rhomboides (Linnaeus, 1766)		●	●	●
			Stenotomus caprinus Jordan & Gilbert, 1882				●

Class	Order	Family	Species	1980-1981	1998-1999	2010-2011	2016-2017
<i>Actinopteri</i>	Scombriformes	Stromateidae	<i>Peprilus paru</i> (Linnaeus, 1758)		●	●	
		Trichiuridae	* <i>Trichiurus lepturus</i> Linnaeus, 1758	●	●	●	●
	Siluriformes	Ariidae	*~ <i>Ariopsis felis</i> (Linnaeus, 1766)	●	●	●	●
			* <i>Bagre marinus</i> (Mitchill, 1815)	●	●	●	●
			*~ <i>Cathorops melanopus</i> (Günther, 1864)	●	●	●	●
	Syngnathiformes	Dactylopteridae	<i>Dactylopterus volitans</i> (Linnaeus, 1758)			●	
		Syngnathidae	<i>Hippocampus erectus</i> Perry, 1810	●	●	●	●
			<i>Hippocampus zosterae</i> Jordan & Gilbert, 1882				●
			<i>Syngnathus floridae</i> (Jordan & Gilbert, 1882)				●
			~ <i>Syngnathus fuscus</i> Storer, 1839			●	
			<i>Syngnathus louisianae</i> Günther, 1870	●	●		●
			~ <i>Syngnathus scovelli</i> (Evermann & Kendall, 1896)	●	●	●	●
	Tetraodontiformes	Diodontidae	<i>Chilomycterus schoepfii</i> (Walbaum, 1792)	●	●	●	●
		Monacanthidae	<i>Aluterus schoepfii</i> (Walbaum, 1792)	●		●	●
			<i>Monacanthus ciliatus</i> (Mitchill, 1818)		●	●	
			<i>Stephanolepis hispida</i> (Linnaeus, 1766)	●	●	●	●
		Ostraciidae	<i>Acanthostracion quadricornis</i> (Linnaeus, 1758)	●	●	●	●
		Tetraodontidae	<i>Lagocephalus laevigatus</i> (Linnaeus, 1766)	●	●	●	
			<i>Sphoeroides greeleyi</i> Gilbert, 1900	●	●	●	●
			<i>Sphoeroides maculatus</i> (Bloch & Schneider, 1801)			●	
			<i>Sphoeroides marmoratus</i> (Lowe, 1838)	●			
			<i>Sphoeroides nephelus</i> (Goode & Bean, 1882)	●	●	●	●
			<i>Sphoeroides pachygaster</i> (Müller & Troschel, 1848)			●	
			<i>Sphoeroides parvus</i> Shipp & Yerger, 1969			●	●
			<i>Sphoeroides spengleri</i> (Bloch, 1785)	●	●	●	●
			~ <i>Sphoeroides testudineus</i> (Linnaeus, 1758)	●	●	●	●



Annex III. List of species of phytobenthos in Laguna de Términos (Ortega, 1995)

	Class	Species
Algae	Cyanophiceae	<i>Aphanocapsa litoralis</i>
		<i>Lyngbya aestuarii</i>
		<i>Lyngbya confervoides</i>
		<i>Lyngbya majuscula</i>
		<i>Microcoleus chthonoplastes</i>
		<i>Phromidium sp</i>
		<i>Calothrix longifila</i>
	Rhodophyceae	<i>stylonema alsidii</i>
		<i>Erythrocladia irregularis</i>
		<i>Erythrocladia sp</i>
		<i>Audouinella hypnea</i>
		<i>Gelidium americanum</i>
		<i>Gelidium pusillum</i>
		<i>Fosliella farinosa</i>
		<i>Fosliella lejolisii</i>
		<i>Jania adhaerens</i>
		<i>Hypnea cervicornis</i>
		<i>Hypnea valentiae</i>
		<i>Hypnea musciformis</i>
		<i>Hypnea spinella</i>
	Rhodophyceae	<i>Hypneocolax stellaris</i>
		<i>Agardhiella ramosissima</i>
		<i>Agardhiella subulata</i>
		<i>Eucheuma gelidium</i>
		<i>Eucheuma inerme</i>
		<i>Eucheuma isiforme</i>
		<i>Gracilaria caudata</i>
		<i>Gracilaria cervicornis</i>
		<i>Gracilaria curtissiae</i>
		<i>Gracilaria cylindrica</i>
		<i>Gracilaria damaecornis</i>
		<i>Gracilaria domingensis</i>
		<i>Gracilaria lemaneiformis</i>
		<i>Gracilaria tikvahiae</i>
		<i>Gracilaria venezuelensis</i>
		<i>Gracilaria verrucosa</i>
		<i>Gracilaria sp</i>
		<i>Gymnogongrus tenuis</i>
		<i>Champia parvula</i>
		<i>Chrysymenia sp</i>
		<i>Anotrichium tenue</i>
		<i>Callithamnion sp</i>

	Class	Species
Algae	Rhodophyceae	<i>Centroceras clavulatum</i> <i>Ceramium diaphanum</i> <i>Ceramium fastigiatum</i> <i>Ceramium flaccidum</i> <i>Spyridia filamentosa</i> <i>Wrangelia argus</i> <i>Caloglossa leprieurii</i> <i>Heterosiphonia crispella</i> <i>Acanthophora spicifera</i> <i>Bostrychia radicans</i> <i>Chondria baileyana</i> <i>Chondria sedifolia</i> <i>Chondria tenuissima</i> <i>Digenea simplex</i> <i>Herposiphonia secunda</i> <i>Laurencia gemmifera</i> <i>Laurencia papillosa</i> <i>Lophocladia trichoclados</i> <i>Polysiphonia ferulacea</i> <i>Polysiphonia sp</i>
		<i>Vaucheria sp</i>
	Xanthophyceae	<i>Ectocarpus? rhodochortonoides</i> <i>Feldmannia elachistaeformis</i> <i>Giffordia indica</i> <i>Hincksia mitchelliae</i> <i>Hincksia sp</i> <i>Dictyota bartayresii</i> <i>Dictyota ciliolata</i> <i>Padina boergesenii</i> <i>Padina gymnospora</i>
		<i>Entocladia ventriculosum</i> <i>Entocladia viridis</i> <i>Phaeophyla dendroides</i> <i>Enteromorpha flexuosa</i> <i>Entheromorpha lingulata</i> <i>Ulva fasciata</i> <i>Cladophoropsis membranacea</i> <i>Chaetomorpha aerea</i> <i>Chaetomorpha antenninna</i> <i>Chaetomorpha linum</i>
		<i>Cladophora vagabunda</i> <i>Rhizoclonium africanum</i> <i>Rhizoclonium kernerii</i>

	Class	Species
Algae	Chlorophyceae	<i>Bryopsis plumosa</i>
		<i>Bryopsis ramulosa</i>
		<i>Codium isthmocladum</i>
		<i>Caulerpa fastigiata</i>
		<i>Caulerpa mexicana</i>
		<i>Caulerpa prolifera v zosterifolia</i>
		<i>Caulerpa prolifera v zosterifolia</i>
		<i>Caulerpa racemosa peltata</i>
		<i>Caulerpa racemosa occidentales</i>
		<i>Caulerpa sertularioides</i>
		<i>Caulerpa sertularioides</i>
		<i>Caulerpa sertularioides longiseta</i>
		<i>Acetabularia crenulata</i>
Seagrass	Monocotyledoneae	<i>Halophyla aschersonii</i>
		<i>Halophyla decipiens</i>
		<i>Thalassia testudinum</i>
		<i>Halodule wrightii</i>
		<i>Syringodium filiforme</i>



REGISTRO DE DATOS DE JAIBA A PIE DE PLAYA

PESCADOS Y MARISCOS DEL CARIBE SA DE CV

Calle Aldama Manzana 90 Lote 2, Col. Banco de Piedra

24100 Sabancuy, Ciudad Del Carmen, Campeche.

FECHA EMISIÓN
2/JUN/2021CLAVE:
FSPR06NO. REVISIÓN
01FECHA REVISIÓN
2/JUNIO/2021PÁGINA
1 DE 1

FOLIO	COMPRADOR <i>Olegario P.</i>			FECHA <i>21/07/2023</i>
LOCALIDAD DE DESEMBARQUE <i>Palo Alto.</i>	ZONA DE PESCA (UBICACIÓN MAPA) <i>El Bajío.</i>	SEMANA <i>3</i>		
PROVEEDOR (ORGANIZACIÓN O PERMISIÓNARIO) <i>Uniser</i>	(NOMBRE EMBARCACIÓN)			
PESCADOR NOMBRE COMPLETO <i>Uniser Hernandez Jimenez.</i>	APODO PESCADOR —	POTENCIA MOTOR —		
AREOS O NASAS #	TRAMPA # <i>80</i>	Eslora (LARGO) —	Manga (ANCHO) —	Puntal (ALTURA) —
KILOS RECIBIDOS (JAIBA Y JAIBÓN) <i>22 kg</i>	KILOS JAIBA —	KILOS JAIBON —		
TIPO DE CARNADA <i>Sardina.</i>	HORAS DEL ARTE DE PESCA EN EL AGUA <i>4 hrs.</i>			
PESO DE CARNADA (K) <i>8kg.</i>				

ESPECIES CAPTURADAS

<i>C. sapidus</i> Azul Cs	<i>C. bocourti</i> Cb
<i>C. rathbunae</i> café, negra Cr	<i>C. danae</i> Cd
<i>C. similis</i> "pata seca" S	<i>C. ornatus</i> Co

INFORMACIÓN DEL HÁBITAT

Tipo de fondo *Verde Socate.*Profundidad *1/2 metro.*

Observaciones generales durante la pesca

Descripción de como se desarrolla la pesca

Fauna asociada

Nombre común	Nombre científico	Cantidad (k)	Se descarta	Se regresa al agua	Consumo propio	Venta
<i>Caracol.</i>		10 Pz	3	7	—	—
<i>Congrio.</i>		5 Pz	—	5	—	—
<i>Trucha.</i>		25 Pz	6	19	—	—
<i>Pez Sapo.</i>		1	—	1	—	—

OBSERVACIONES. Ejemplo: observación ocasional de otras especies como Arenareus spp. * Datum NAD27 (North American Datum 1927) o WGS84 (World Geodetic System 1984)

ELABORÓ:

Orlando Garrido

SUPERVISO:

Joa R Z

FSPR06 rev 01