

## Productivity Susceptibility Analysis (PSA) of bycatch in Yucatan Peninsula blue crab *Callinectes sapidus* trap fishery based on observed data

Fisheries Improvement Project Mexico Yucatan Peninsula blue crab-dipnet/pot/trap



September, 2020

Minerva N. Alonso Alemán<sup>1</sup>, B. Gabriela Mc Lean<sup>2</sup>; Ernesto Godelman<sup>2</sup>

<sup>1</sup> CeDePesca México, Mérida Yucatán. E-mail: [minerva.alonso@cedepesca.net](mailto:minerva.alonso@cedepesca.net)

<sup>2</sup>Centro Desarrollo y Pesca Sustentable (CeDePesca). Rondeau 361. (B7603BDG). Mar del Plata. Buenos Aires. Argentina.

## Content

Summary .....	3
Acronyms .....	3
Introduction .....	3
Data.....	3
Methodology .....	4
Discussion .....	8
<i>Spherooides testudineus</i> (Linnaeus, 1758).....	8
<i>Ariopsis melanopus; Ariopsis felis, Bagre marinus</i> .....	10
<i>Archosargus rhomboidalis</i> .....	12
<i>Achirus lineatus</i> .....	14
<i>Eucinostomus gula</i> .....	16
<i>Diapterus rhombeus</i> .....	18
<i>Clibanarius vittatus</i> .....	20
<i>Callinectes similis</i> .....	22
<i>Callinectes rathbunae</i> .....	24
<i>Hepatus epheliticus</i> .....	26
<i>Libinia dubia</i> .....	28
<i>Menippe mercenaria</i> .....	30
Results .....	32
Conclusions.....	33
Literature cited .....	34



## Summary

This work applies the risk assessment method known as Productivity and Susceptibility Analysis (PSA) to the crustaceans: lesser blue crab, locally known as “pata seca” (*Callinectes similis*), sharptooth swimcrab (*Callinectes rathbunae*), hermit crab (*Clibanarius vittatus*), calico box crab (*Hepathus epheliticus*), stone crab (*Menippe mercenaria*), lognose spider crab (*Libinia dubia*), and to the fish: checkered puffer (*Sphoeroides testudineus*), hardhead catfish (*Ariopsis felis*), Western Atlantic seabream (*Archosargus rhomboidalis*), lined sole (*Achirus lineatus*), jenny mojarra (*Eucinostomus gula*) & mojarra (*Diapterus rhombeus*) species that have appeared as part of the accompanying fauna in the blue crab ring trap fishery in Isla Aguada and Sabancuy (Laguna de Términos) in Campeche.

## Acronyms

<b>CSIRO</b>	Commonwealth Scientific and Industrial Research Organización
<b>ERAEF</b>	Environmental Risk Assessment for the Effects of Fishing
<b>GMFMC</b>	Gulf of Mexico Fishery Management Council
<b>MSC</b>	Marine Stewardship Council
<b>PSA</b>	Productivity-Susceptibility Analysis
<b>RBF</b>	Risk-Based Framework

## Introduction

Productivity-Susceptibility Analysis (PSA) is a semi-quantitative method used to assess the vulnerability of species associated within data deficient fisheries. The method has been refined by the Commonwealth Scientific and Industrial Research Organization (CSIRO) of Australia, based on the method developed by Stobutzki et al. (2001) to evaluate the probability of sustainability of the species of the companion fauna in a fishery. The PSA is part of a broader hierarchical methodology developed by CSIRO, called the Environmental Risk Assessment for the Effects of Fishing (ERAEF), which can be applied to species, habitats or communities with low fishing impact (Hobday et al., 2004; Hobday et al., 2007) and has been used to assess all fisheries managed by the Australian Fisheries Management Authority (Smith et al., 2007) and it is part of the Marine Stewardship Council (MSC) framework.

## Data

A preliminary list of species that have appeared within the crab traps used in the fishery was made in the samplings carried out between October 2018 and December 2019. The samplings were carried out at the Reception Center located in Isla Aguada, Campeche where fishermen

take their catches in order to sell them to the organization. Many of the organisms observed were returned to the sea alive. Not all the accompanying fauna reaches the reception center because fishermen frequently release live species caught incidentally and of no commercial interest in the fishing area. This list of species was completed with the results of interviews with fishermen carried out during the same period. The identification of the species was carried out from Ayala-Perez, et al (2015), Perry and Larsen, 2004, Raz-Guzman, 1999, the FAO guide prepared by Fischer et al 1995) and the one prepared by Carpenter, (2002). A total of 12 species were identified as bycatch.

## Methodology

In order to determine the risk level for each species, a bibliographic search was carried out in order to get the information available on the species in the region. When species-specific information was not available, information from similar species within the same genus or family was used. When this was not possible, a precautionary approach was considered by assigning the highest risk value.


Productivity and Susceptibility attributes were scored based in tables 1 & 2, and on the information gathered. Trophic level attributes were obtained from Fishbase (2020), Sealifebase (2020) or any other detailed information for crustaceans.

The score for each component of the PSA was recorded in the “MSC RBF Worksheet”, and rationales were included when no specific information was found.

The PSA (Figure 1) is based on the assumption that the potential risk of fishing to a species that is directly or indirectly impacted depends on two main components (Stobutzki et al., 2001; Stobutzki et al., 2002; Hobday et al., 2007; Patrick et al., 2010; Hobday et al., 2011; Clarke et al., 2018):

- ❖ Its natural productivity, related to the recovery capacity of the species. This attribute is associated with the biology and ecology of the species..
- ❖ Its susceptibility to being fished, which reflects how exposed it is to fishing activities. This component is related to the characteristics of both the fishing gear and the spatial distribution of the fishery in relation to its natural distribution in the area.

The potential risk associated with the productivity or susceptibility of a particular species was determined with the tools provided by MSC (v 2.1) for applying the RBF.



**Figure 1. PSA graph with productivity and susceptibility axes, and the risk space divided into regions. If the vulnerability is  $<1.80$ , the risk is low; if the vulnerability is between  $1.80$  and  $2.20$ , the risk is medium; and if the vulnerability is  $>2.20$ , the risk is high. The graph has been modified from Stobutzki et al. (2001b), inverting the productivity scale that starts with 3 (high productivity) instead of 1 (low productivity).**

The PSA considers eight productivity attributes: average age at maturity, average size at maturity, average maximum age, average maximum size, fecundity, reproductive strategy, trophic level and density dependence (only for invertebrates). The average maximum age and average maximum size are not used for the evaluation of invertebrates. For each of these attributes there are ranges associated with different levels of productivity as presented in Table 1. If the level of productivity associated with the attribute is high, a score of 1 (low risk) is given; if it is medium, 2 (medium risk); and if it is low, 3 (high risk).

**Table 1. PSA productivity attributes and scores (from MSC V 2.0)**

Productivity attribute	High productivity (Low risk, score = 1)	Medium productivity (medium risk, score = 2)	Low productivity (high risk, score = 3)
Average age at maturity	<5 years	5-15 years	>15 years
Average maximum age	<10 years	10-25 years	>25 years
Fecundity	>20,000 eggs per year	100-20,000 eggs per year	<100 eggs per year
Average maximum size (not to be used when scoring invertebrate species)	<100 cm	100-300 cm	>300 cm
Average size at maturity (not to be used when scoring invertebrate species)	<40 cm	40-200 cm	>200 cm
Reproductive strategy	Broadcast spawner	Demersal egg layer	Live bearer
Trophic Level	<2.75	2.75-3.25	>3.25
Density dependence !! (to be used when scoring invertebrate species only)	Compensatory dynamics at low population size demonstrated or likely.	No dependatory or compensatory dynamics demonstrated or likely.	Depensatory dynamics at low population sizes (Allee effects) demonstrated or likely.

In the case of susceptibility, there are four attributes: availability, which refers to the portion of the horizontal distribution of the species that coincides with the areas in which the fishery takes place; encounterability, which refers to the vertical overlap between the usual position of the fishing gear and the species in the water column; selectivity of fishing gear, and post-capture mortality (Hobday, 2007; Smith et al., 2007). Selectivity of the fishing gear is evaluated based on two criteria: the relationship between the size of immature individuals and the probability of their capture; and the possibility that these immature individuals may escape the fishing gear alive or may evade it.



Each attribute receives a score of 1, 2, or 3, depending on whether it is associated with low (low risk), medium (medium risk), or high (high risk) susceptibility, in accordance to criteria shown in the Table 2.

**TABLE 2. Susceptibility attributes (MSC version 2.1)**

Susceptibility attribute	High productivity (Low risk, score = 1)	Medium productivity (medium risk, score = 2)	Low productivity (high risk, score = 3)
Areal overlap (availability) Overlap of the fishing effort with a species concentration of the stock	<10% overlap	10-30% overlap	>30% overlap
Encounterability The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear	Low overlap with fishing gear (low encounterability).	Medium overlap with fishing gear.	High overlap with fishing gear (high encounterability).  Default score for target species (Principle 1).
Selectivity of gear type Potential of the gear to retain species	a Individuals < size at maturity are rarely caught.	a Individuals < size at maturity are regularly caught.	a Individuals < size at maturity are frequently caught.
	b Individuals < size at maturity can escape or avoid gear.	b Individuals < half the size at maturity can escape or avoid gear.	b Individuals < half the size at maturity are retained by gear.
Post-capture mortality (PCM) The chance that, if captured, a species would be released and that it would be in a condition permitting subsequent survival	Evidence of majority released post-capture and survival.	Evidence of some released post-capture and survival.	Retained species or majority dead when released.  Default score for retained species (Principle 1 or Principle 2).

## Discussion

Specific information for each species, references and information of each score is detailed as follows. Most of the scores were evaluated from bibliographic information. In some cases where specific information was not found, genus or family information was taken into consideration as reference. In cases where special criteria were used, a brief analysis is included. Fish trophic level was taken from Fishbase in all cases (Froese & Pauly, 2019).

### *Sphoeroides testudineus* (Linnaeus, 1758)

Botete o Checkered puffer

Inhabits shallow waters up to 10 m in marine and brackish habitats (Shao et al, 2014). It is distributed from Rhode Island, USA to southern Brazil (Wester Atlantic to Southwest Atlantic). Commonly found in bays, tidal creeks and protected coastal waters, especially on seagrass beds, and in brackish water. Rare or absent on coral reefs. Does not form schools, but may form huge aggregates. Hides in the sand when frightened (Lieske & Myers, 1994). Feeds mainly on bivalves, gastropods, foraminiferans and several other benthic invertebrates specially crustaceans, which it crushes with its powerful teeth (Keith et al, 2000). To ward off predators, it inflates itself like a balloon. Highly toxic. (Pauly, 1991). Spawning season is from late spring to early fall at Biscayne Bay, Florida. Mean length at first maturity is over 13 cm TL. (Pauly, 1991)

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	< 1.53 y (Bonilla et al, 2012)	1
Average maximum age	1.53 /y (Bonilla et al 2012)	1
Fecundity	1.146 eggs/body weight Targett, T.E., 1979; 59 087 and 367 022 oocytes (mean 176 456) per spawning batch for <i>P. nephelus</i> (Peniche-Pérez, et al, 2019)	1
Average maximum size	38.8 cm (Claro, 1994)	1
Average size at maturity	10.8 cm or 13 cm. (Pauly, 1991)	1
Reproductive strategy	Batch spawning (Peniche-Pérez, et al, 2019), since it is a demersal fish it is considered	2
Trophic level	3.4	3
Density dependence (only for invertebrates)		



Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort with a species concentration of the stock	This species is found along the whole Laguna de Términos (Mallard et al, 1981) and it is distributed from Rhode Island to Brazil. Fishing effort is concentrated, it is considered as a precautionary approach 10-30% overlap	2
Encounterability The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear	Since it is a demersal fish, it is considered a high overlap with fishing gear.	3
Ring traps Selectivity of gear type Potential of the gear to retain species	Individuals < size at maturity are regularly caught (from interviews with fishermen)	2
Post-capture mortality (Considers fishery under assessment)	Individuals < half the size at maturity can escape or avoid gear & Fishermen can free them alive since it has no commercial interest.	2



*Ariopsis melanopus; Ariopsis felis, Bagre marinus*

Hardhead sea catfish, locally known as bagre

Sea catfishes occur in marine, brackish, and fresh waters of warm-temperate and tropical regions. The marine representatives are mostly confined to the coastlines of the continent and continental islands such as Trinidad (only a single species has been reported from Cuba, for example). They may be locally abundant in the turbid waters of certain habitats, particularly large river estuaries and mangrove-lined lagoons. A few species may reach depths of 100 m or deeper. Sea catfishes' diets range from omnivorous, including detritus, to strongly carnivorous, including large bony fishes and crustaceans. Reproduction is highly specialized: the males incubate the eggs and vitelline young in their mouths. The sea catfishes include several species of high economic value. FAO statistics report landings ranging from 14 885 to 26 630 t from 1995 to 1999. They are captured with a variety of gear, including bottom trawls, longlines, seines, cast nets, traps, and hook-and-line. The flesh is usually of good quality, but the sharp and serrated dorsal- and pectoral-fin spines can inflict painful wounds. (Acero, A., 2002).

Wide distribution from the US to Venezuela, depending on the species. In the U.S. Gulf of Mexico, where marine catfishes remain common and can be found in such high local abundances that some fisheries reportedly avoid areas due to unwanted marine catfish bycatch (Armstrong et al. 1996). Winemiller & Rose (1992) consider this fish exhibit an equilibrium life history strategy. Armstrong et al. (1996) estimated maturity at around age 5.

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	5 years (Armstrong, et al, 1996).	2 As a precautionary approach, given the long life span of the species, a score of 2 is considered instead of 1.
Average maximum age	24 years (Flinn, et al, 2019)	2
Fecundity	15 eggs found in mouth, may bear up to 43, (Segura-Bertolini, 2011)	3
Average maximum size (not for invertebrate)	70 cm (Fishbase)	1
Average size at maturity (not for invertebrate)	13 cm (Fishbase)	1
Reproductive strategy	Bearers/external brooders (Fishbase)	3
Trophic level	3.2	3
Density dependence (only for invertebrates)		

Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort	This species has a wide distribution in Laguna de	1

with a species concentration of the stock	Términos and it is important in terms of abundance (Ayala-Pérez, et al 2012). Fishing effort is concentrated, it is considered as a precautionary approach 10-30% overlap	
<p>Encounterability</p> <p>The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear</p>	This group of species feed on the seabed, having a strong interaction with fishing gear.	3
<p>Ring traps</p> <p>Selectivity of gear type</p> <p>Potential of the gear to retain species</p>	Individuals < 13 cm (average size at maturity) are rarely caught. It is not clear if they can escape or just avoid gear. As a precautionary approach it is rated as 3.	3
<p>Post-capture mortality (Considers fishery under assessment)</p>	Once in the boat, the fish dies. Fishermen release them since there is no commercial interest, although these are sometimes kept. They are very common in the area, a score of 3 was chosen as a precautionary approach.	3

*Archosargus rhomboidalis*

Commonly found over mud bottoms in mangrove sloughs and on vegetated sand bottoms, sometimes in brackish water and occasionally also in coral reef areas near mangroves. Feeds on benthic invertebrates (small bivalves, crustaceans), as well as on plant material. Distribution USA and northeastern Gulf of Mexico to Argentina.

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	< 2 years (deduced from Av Max Age)	1
Average maximum age	2 years (Chavance, et al 1986)	1
Fecundity	485 eggs (Chavance et al 1986)	2
Average maximum size (not for invertebrate)	33 cm (Robins & Ray, 1986)	1
Average size at maturity (not for invertebrate)	Lm 8 cm (Fishbase)	1
Reproductive strategy	Non guarders open water/substratum egg scatterers (Chavance et al 1986)	2
Trophic level	2.9	2
Density dependence (only for invertebrates)		

Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort with a species concentration of the stock	This is one of the most abundant species in Laguna de Términos. (Ayala-Pérez, 2003). As a precautionary approach it is scored 2.	2
Encounterability The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear	Commonly found over mud bottoms in mangrove sloughs and on vegetated sand bottoms	3



<p>Ring traps</p> <p>Selectivity of gear type</p> <p>Potential of the gear to retain species</p>	<p>Biological Monitoring Program suggests individuals &lt; size at maturity (8 cm) are rarely caught. No information exists if Individuals &lt; size at maturity can avoid or escape gear.</p>	<p>3</p>
<p>Post-capture mortality (Considers fishery under assessment)</p>	<p>Default score for retained species as fishermen use it as bait</p>	<p>3</p>

*Achirus lineatus*

Lined sole

It is distributed along the Western Atlantic, from Florida and Northern Mexico to northern Argentina. Occurs mainly in brackish or hyper-saline lagoons, on sandy-muddy bottoms of estuaries and in the littoral zone from 1 to 20m depths. Its growth rate is relatively slow (Keith, et al. 2000).

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	Deducted from average maximum age, as a precautionary approach	2
Average maximum age	<i>Trinectes sp.</i> 6 years (Robins & Ray, 1986)	1
Fecundity		3
Average maximum size (not for invertebrate)	<i>Achirus lineatus</i> 33.1 cm; (Ayala-Pérez, et al 2015)	1
Average size at maturity (not for invertebrate)	< 33.1 cm (Ayala-Pérez, et al 2015)	1
Reproductive strategy	<i>Achirus</i> is dioicus; (Ayala-Pérez, et al 2015)	3
Trophic level	3.5 fishbase	3
Density dependence (only for invertebrates)		

Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort with a species concentration of the stock	It has a wide distribution in Laguna de Términos and in the Western Atlantic (Ayala-Pérez et al, 2015)	1
Encounterability The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear	Lives on sandy-muddy bottoms (Keith, et al 2000)	3
Ring traps Selectivity of gear type	Biological Monitoring Program suggests individuals < size at maturity (33.1 cm) may be frequently	3





<p>Potential of the gear to retain species</p>	<p>caught. No information exists if Individuals &lt; size at maturity can avoid or escape gear but this seems not probable since maturity size. As a precautionary approach it is considered the default score 3.</p>	
<p>Post-capture mortality (Considers fishery under assessment)</p>	<p>Fishermen release them since there is no commercial interest, but once in the boat it may die, since this fish does not represent a threat for fishermen they focus their attention in any spine-fish. As a precautionary approach it is considered the default score 3.</p>	<p>3</p>

*Eucinostomus gula*

Jenny mojarra

Gerreidae family are chiefly marine. In brackish water, it is encountered occasionally; rarely, in freshwater. Very protractile mouth. Head scaly but with smooth upper surface. Dorsal and anal fins with a sheath of scales along base. Gill membranes not united to isthmus. Deeply forked tail; 24 vertebrae. Maximum length 35 cm, attained in *Gerres filamentosus*. Small silvery fishes with highly protrusible mouth. They feed by sorting benthic invertebrates from sand. Foodfishes. Assumed to be nonguarders. Most of the adult species occur in coastal lagoons with sandy or muddy bottoms bordered by mangroves; however, they occasionally enter river mouths. Juveniles enter estuaries until they reach maturity; spawning occurs at sea throughout the seasons. (Fishbase).

The mojarras were defined as second-order consumers, feeding on microcrustaceans (amphipods, copepods, tanaidaceous, ostracods) and significant amounts of detritus with variations in the proportion and frequency of different types of food according to their ontogeny and food availability (Chi-Espínola, 2018). Mojarras are omnivorous and euriphagous fishes, eating a wide spectrum of items, mainly small invertebrates as bivalves, gastropods, ostracods, copepods, polychaetes (Arenas-Granados & Acero, 1992).

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	No information found. Default score selected.	3
Average maximum age	No information found. Default score selected.	3
Fecundity	111 eggs (Mexicano-Cántora)	3
Average maximum size (not for invertebrate)	25.5 cm (Amador, et al. 2015)	1
Average size at maturity (not for invertebrate)	11 cm (Amador, et al. 2015)	1
Reproductive strategy	Broadcast spawner	1
Trophic level	2.7 (Fishbase)	2
Density dependence (only for invertebrates)		

Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort with a species concentration of the stock	Broad distribution	1

<p>Encounterability</p> <p>The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear</p>	<p>Inhabits shallow waters, being especially abundant over mud bottoms in mangrove-lined lagoons or creeks; larger individuals may also occur on vegetated sand grounds in marine areas.</p>	<p>3</p>
<p>Ring traps</p> <p>Selectivity of gear type</p> <p>Potential of the gear to retain species</p>	<p>Biological Monitoring Program suggests individuals &lt; size at maturity (11 cm) may be rarely caught. No information exists if Individuals &lt; size at maturity. As a precautionary approach it is considered the default score 3.</p>	<p>3</p>
<p>Post-capture mortality (Considers fishery under assessment)</p>	<p>Default score for retained species as fishermen use it as bait.</p>	<p>3</p>

*Diapterus rhombeus*

Mojarra

Distributed from Gulf of Mexico, Central America and the Antilles to Brazil (Cervigón, F., 1993). Common in mangrove-lined lagoons, found over shallow mud and sand grounds in marine areas. Juveniles common in hypersaline lagoons and in brackish water (Cervigón, 1992). Lives from 9-70 m depth. Annual catches were estimated from 100-200 t (Claro, 1994).

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	No information found. Default score selected.	3
Average maximum age	No information found. Default score selected.	3
Fecundity	No information found. Default score selected.	3
Average maximum size (not for invertebrate)	40 cm (Randall y Vergara, 1978 in Ayala-Pérez et al, 2015)	1
Average size at maturity (not for invertebrate)	9 cm (Rodríguez da Costa, et al 2012)	1
Reproductive strategy	Its spawning most likely occurs in adjacent coastal waters, (Elliff et al, 2013). Previous observations made on <i>D. rhombeus</i> revealed that adults typically spawn in areas deeper than 10 m, whereas juveniles use the shallow waters of estuaries, bays and mangroves. Chaves et al (1998) suggested that smaller individuals (i.e., <150 mm TL) present evidence of previous spawning events. Presents breeding aggregation & a unique nest-clearing behavior, the eggs potentially released over the cleared surfaces were nearly undistinguishable from sediment particles. (Reis-Filho, 2018)	2
Trophic level	3.0 (Fishbase)	2
Density dependence (only for invertebrates)		

Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort with a species concentration of the stock	Broad distribution	1
Encounterability  The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear	Between 9 & 70 m (Gines & Cervigón, 1967). It is common to find it in coastal mangrove areas of coastal lagoons with muddy bottoms and in deep sea areas Sandy. (Froese y Pauly, 2009)	3
Ring traps  Selectivity of gear type  Potential of the gear to retain species	Biological Monitoring Program suggests individuals < size at maturity (9 cm) may be rarely caught. No information exists if Individuals < size at maturity. As a precautionary approach it is considered the default score 3.	3
Post-capture mortality (Considers fishery under assessment)	Default score for retained species as fishermen use it as bait.	3

*Clibanarius vittatus*  
Cangrejo hermitaño

Adult crabs are commonly found inhabiting gastropod shells at least 10 cm long (Ruppert & Fox 1988). Thin stripe hermits are considered to employ an opportunistic trophic mode, feeding on a variety of plant and animal material. Diet studies involving gut content analysis revealed that crabs consumed 40% scavenged material, 40% detritus, and 20% substratum (Williams 1984).

Information on the predators of *C. vittatus* is scarce, but crabs are likely preyed upon by large benthic-feeding fishes and other crustaceans. Eggs and larvae are consumed by a variety of organisms, including some species that are considered to be commensals of the parent crab (Williams 1984).

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	No information found. Default score selected.	3
Average maximum age	No information found. Default score selected.	3
Fecundity	180,000 eggs (Turra & Leite, 2001)	1
Average maximum size (not for invertebrate)		
Average size at maturity (not for invertebrate)		
Reproductive strategy	Females carries eggs (sponge) Broadcast spawner	1
Trophic level	No information found. Default score selected.	3
Density dependence (only for invertebrates)	No dependantory or compensatory dynamics demonstrated or likely	2

Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort with a species concentration of the stock	Wide distribution of the species (Crab data base)	1



<p>Encounterability</p> <p>The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear</p>	<p>Since they live on the bottom, there is a high probability of encounter</p>	<p>3</p>
<p>Ring traps</p> <p>Selectivity of gear type</p> <p>Potential of the gear to retain species</p>	<p>Individuals below size at maturity are regularly caught (5-50%)</p>	<p>2</p>
<p>Post-capture mortality (Considers fishery under assessment)</p>	<p>Evidence of majority released post-capture and survival</p>	<p>1</p>

*Callinectes similis*  
Lesser blue crab

Adult *Callinectes similis* occur in the oceanic littoral zone in salinities above 15‰ at a depth of 100 meters along the eastern seaboard from Delaware Bay to Key West, Florida and southern coasts of the United States, to the Yucatan, Colombia and northern Jamaica (Williams 1984, Piller et al. 1995). It is distributed throughout the lagoon independently of a particular type of habitat (Sánchez & Raz-Guzman, 1997). *Callinectes similis* was most frequently (92%) collected in salinities above 20‰, in areas with macroalgae, sea grasses, and submerged aquatic vegetation (70%) and on unvegetated soft substrates (30%) (Sánchez & Raz-Guzman, 1997). *C. similis* is omnivorous; its diet includes detritus and invertebrates, and carrion (Britton & Morton, 1989). La jaiba *Callinectes similis* da soporte a la pesquería de plataforma. (Raz-Guzmán, 2005)

Estudio de caso: crustáceos de la laguna de Términos. En: Diversidad de especies: estudio de caso. 275-278.pp.

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	As of <i>Callinectes sapidus</i> , 12-18 months of age (Rosas-Correa and Jesus-Navarrete 2008) (Estrada-Valencia, 1999) (Fischer & Wolff, 2006) (ADW, 2016)	1
Average maximum age	As of <i>Callinectes sapidus</i> , 4.5 years (Rosas-Correa and Jesus-Navarrete 2008)	1
Fecundity	277,886 ±136,270 (Chazaro-Olvera et al, 2000); 125,734 to 986,393 eggs (García-Montes, et al., 1987)	1
Average maximum size (not for invertebrate)		
Average size at maturity (not for invertebrate)		
Reproductive strategy	Broadcast spawner	1
Trophic level	Unknown, 3.3 (as <i>C. sapidus</i> )	3
Density dependence (only for invertebrates)	No density dependence or compensatory dynamics demonstrated or likely	2

Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort	Distributed throughout Laguna de Términos, found	3

<p>with a species concentration of the stock</p>	<p>on all sorts of substrates along a wide range of salinity (Sánchez &amp; Ráz-Guzmán, 1997), prefers salinities over 15‰. (Piller, et.al. 1995) Broad distribution, considered a precautionary approach since not much information found. Common in the Campeche area, specially in the coastal platform, sustains the platform fishery. (Raz-Guzmán, 2010)</p>	
<p>Encounterability The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear</p>	<p>Since they live on the bottom, there is a high probability of encounter</p>	<p>3</p>
<p>Ring traps Selectivity of gear type Potential of the gear to retain species</p>	<p>Individuals below size at maturity are regularly caught (5-50%)</p>	<p>2</p>
<p>Post-capture mortality (Considers fishery under assessment)</p>	<p>FIP has evidence that fishermen release them, but when catches are scarce, they keep it, as a precautionary approach it is scored as default.</p>	<p>3</p>

### *Callinectes rathbunae*

Where no specific information for *C. rathbunae*, information of the genera is included.

*Callinectes rathbunae* was collected from all types of habitats along the margins of the NW, W, C-SE, and S-SW areas of the Laguna de Términos lagoon throughout wide ranges of salinity (3-34‰), on sand (5-99%), on mud (1-94%), as well as on habitats with and without submerged aquatic vegetation (Sánchez & Raz-Guzman, 1997). However, it is not frequently caught by fishermen, since they are looking for blue crabs (local fishermen information).

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	As of <i>Callinectes sapidus</i> , 12-18 months of age (Rosas-Correa and Jesus-Navarrete 2008) (Estrada-Valencia, 1999) (Fischer & Wolff, 2006) (ADW, 2016)	1
Average maximum age	As of <i>Callinectes sapidus</i> , 4.5 years (Rosas-Correa and Jesus-Navarrete 2008)	1
Fecundity	As of <i>Callinectes sapidus</i> , 872,000 eggs/year (Estrada-Valencia, 1999; ADW, 2016)	1
Average maximum size (not for invertebrate)		
Average size at maturity (not for invertebrate)		
Reproductive strategy	Brooder	2
Trophic level	As of <i>Callinectes sapidus</i> , 3.3 (Morales-Zarate, et al. 2004); 3.73 (Mancinelli, et al, 2016)	3
Density dependence (only for invertebrates)	As of <i>Callinectes sapidus</i> , No dependant or compensatory dynamics demonstrated or likely	2

Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort with a species concentration of the stock	Broad distribution. Frequently found in freshwater mouths or close to them. (Sánchez & Raz-Guzman, 1997)	1

<p>Encounterability</p> <p>The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear</p>	<p>Since it inhabits bottom a precautionary approach is scored.</p>	<p>3</p>
<p>Ring traps</p> <p>Selectivity of gear type</p> <p>Potential of the gear to retain species</p>	<p>No information found. Default score</p>	<p>3</p>
<p>Post-capture mortality (Considers fishery under assessment)</p>	<p>Since species is kept when caught, default score for retained species</p>	<p>3</p>

*Hepatus epheliticus*

Cangrejo fuente o calico box crab

*Hepatus epheliticus*. is an abundant crab commonly found in shallow and continental shelf waters of the Atlantic coast of USA, Mexico (Negreiros-Fransozo, 2008; Hernáez, 2012) Cuba, Jamaica and Dominican Republic (Williams, 1984). It lives buried in sandy substrate between depths of 2 to 91 m (Franks et al., 1972). Hernáez (2012) suggests that ovigerous females may migrate to other areas away from the fishing area in the Campeche Bank or since baited traps were used in their study, that ovigerous females do not enter in them as recorded by other authors. This crab is caught to be used as bait for the octopus fishery in the Campeche Bank, Mexico (Hernáez, 2012).

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	No information found. Default score selected.	3
Average maximum age	Longevity for <i>H. pudibundus</i> was estimated at 1.91 and 2.40 years for males and females, respectively (Fernandes-Miazaki, 2018)	1
Fecundity	75,615 +/- 30,120 eggs/female/spawn. (Reigada, 1995)	1
Average maximum size (not for invertebrate)		
Average size at maturity (not for invertebrate)		
Reproductive strategy	Members of the order Decapoda are mostly gonochoric. Mating behavior: Precopulatory courtship ritual is common (through olfactory and tactile cues); usually indirect sperm transfer (Ruppert et al, 2004). Broadcast spawner (De Lima, 2014).	1
Trophic level	2.75 to 3.25 (Mantellano, 1997 in Seafood Watch, 2019)	2
Density dependence (only for invertebrates)	No evidence for compensatory or dependatory dynamics	2



Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
<p>Aerial overlap (availability) Overlap of the fishing effort with a species concentration of the stock</p>	<p>It is an abundant subtidal crab that lives buried in sandy substrate between depths of 2 to 91 m (Franks, et al., 1972) lives in shallow water in the western Atlantic Ocean from the Chesapeake Bay to the Dominican Republic (Felder, et al, 2009), common in the Campeche area (Raz-Guzmán, 2010)</p>	<p>1</p>
<p>Encounterability The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear</p>	<p>This is a benthic organism (sealifease) commonly found in the ringtraps. The fishing gear is directed to a very similar species so a Default score is considered.</p>	<p>3</p>
<p>Ring traps Selectivity of gear type Potential of the gear to retain species</p>	<p>Default score. Species is not targeted but caught incidentally regularly. As a precautionary approach it is scored 3. The absence of ovigerous females suggests that they did not enter into the traps during embryogenesis. (Hernández, 2012)</p>	<p>3</p>
<p>Post-capture mortality (Considers fishery under assessment)</p>	<p>Evidence of some released post-capture &amp; survival from local fishermen, but when octopus fishery (aug-mid dec) sometimes used as bait for this fishery.</p>	<p>2</p>

*Libinia dubia*

Longnose Spider Crab or maxkil

This crab is commonly referred to as decorator crabs. It is found in a variety of coastal and estuarine habitats to approximately 50 m depth (Williams 1984). The native range of *L. dubia* extends from Cape Cod to southern Texas, Bahamas, Cuba and Yucatan Peninsula (www.inaturalist.org). In Florida, *L. dubia* has been documented as a common inhabitant of Florida Bay (Tabb & Manning 1961), and the most prevalent spider crab in Tampa Bay (Dragovich & Kelly 1964). In Laguna de Términos it was distributed within a narrow salinity range 24-33‰ (Sánchez & Raz-Guzman, 1997). In Indian River Lagoon adults may inhabit more open sandy-bottom areas (Tunberg & Reed, 2004 in <http://naturalhistory2.si.edu>). Commonly used as bait for the Yucatan Peninsula fishery (Bravo-Calderón, et al. 2016). Local populations of *L. dubia* have declined due to overexploitation (CONABIO, 2010) as it is used as bait for the octopus fishery.


Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	About 12 months for <i>L. ferreirae</i> (Gonçalves, et al. 2020)	1
Average maximum age	For <i>Libinia ferreirae</i> 4.68 to 5.91 years (Gonçalves, et al. 2020)	1
Fecundity	20,637 ( $\pm$ SD 12,683.33), with an 81% ( $\pm$ SD 4.62) viability until hatching Carmona-Osalde & Rodríguez-Serna, 2012)	2
Average maximum size (not for invertebrate)		
Average size at maturity (not for invertebrate)		
Reproductive strategy	Broadcast spawner	1
Trophic level	No specific information found. Estimated values after Morales-Zárate, (et al., 2004), considered as “Crab” trophic level 3.30	2
Density dependence (only for invertebrates)	No information found. Default score	3

Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort	This species has a broad distribution from Massachusetts, USA to Cuba and Peninsula de	1

with a species concentration of the stock	Yucatan. It is preferentially distributed in carbonated environments (Soto, 1980). FIP effort overlaps in <10% of its distribution.	
Encounterability  The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear	No information found. Default score	3
Ring traps  Selectivity of gear type  Potential of the gear to retain species	No information found. Default score	3
Post-capture mortality (Considers fishery under assessment)	Retained species when octopus fishery is open	3

## Menippe mercenaria

There is a commercial fishery for Stone crab in the state of Campeche, however, according to NOM-045-SAG / PESC-2015, no place in the Laguna de Terminos appears within the landing areas, therefore its capture is not allowed. . However, with some frequency, some fishermen are observed with the characteristic claw, due to its high commercial value and the custom of self-consumption. The distribution of the Stone crab in the Laguna de Terminos is observed associated with the distribution of underwater vegetation (Sánchez & Raz-Guzmán, 1997) and is reflected in the figure below.



**Fig. *Menippe mercenaria* distribution in Laguna de Términos after Sánchez & Raz-Guzmán, 1997)**

Productivity Attribute	Relevant information	Score (1 low risk, 2 medium risk, 3 high risk)
Average age at maturity	2 years (GMFMC, 1979), <5 years (Crowley, 2018)	1
Average maximum age	2 years (GMFMC, 1979); < 10 years (Gerhart & Bert, 2008)	1
Fecundity	The number of eggs per batch depends on carapace widths, increased with increasing size-class. Number of eggs per batch was found between 40,438 & 836,644 eggs. Additionally, batch fecundity was 39% lower for crabs with no claws, indicating that claw removal by the fishery negatively affects reproductive output. (Crowley, et al, 2019;	1

	Carmona-Osalde & M. Rodríguez-Serna, 2012)	
Average maximum size (not for invertebrate)		
Average size at maturity (not for invertebrate)		
Reproductive strategy	Demersal egg brooder (Crowley, 2019)	2
Trophic level	2.75 to 3.25 (Abeels, et al 2012; Thera et al, 2014)	2
Density dependence (only for invertebrates)	No depensatory or compensatory dynamics demonstrated or likely	2

Susceptibility Attribute	Relevant Information	Score (1 low risk, 2 medium risk, 3 high risk)
Aerial overlap (availability) Overlap of the fishing effort with a species concentration of the stock	Broad distribution, fishery overlaps between 10-30% of its habitat in the coastal lagoon.	2
Encounterability. The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear	Benthic, burrows in mudflats, rocks offshore reef areas or grass clumps. (Froese & Pauly, 2019) Targeted species so default score is used.	3
Ring traps Selectivity of gear type Potential of the gear to retain species	Default. Since stone crabs are a targeted species.	3
Post-capture mortality (Considers fishery under assessment)	Stone crabs although illegal, are a targeted species but released alive after removal of the claws. (Gandy, 2016 & fishermen information)	1

## Results

The final table that includes the scores for each attribute is presented below. According to this results, the MSC score for PI 2.2.1 would be 90 points (Unconditional Pass).

Family name	Scientific name	Common name	Fishery descriptor	Productivity Scores [1-3]								
				Average age at maturity	Average max age	Fecundity	Average max size	Average size at Maturity	Reproductive strategy	Trophic level	Density Dependence	Total Productivity (average)
Tetraodontidae	Sphoeroides testudineus	Botete	Ringtraps	1	1	1	1	1	1	3	3	1.29
Ariidae	Ariopsis felis	Bagre	Ringtraps	2	2	3	1	1	3	3		2.14
Sparidae	Archosargus rhomboidalis	Postá	Ringtraps	1	1	2	1	1	2	2		1.43
Achiridae	Achirus lineatus	Lenguado	Ringtraps	2	1	3	1	1	3	3		2.00
Diogenidae	Clibanarius vittatus	Cangrejo hermitaño	Ringtraps	3	3	1			1	3	2	2.17
Portunidae	Callinectes similis	Jaiba pata seca	Ringtraps	1	1	1			1	3	2	1.50
Portunidae	Callinectes rathbunae	Jaiba negra o café	Ringtraps	1	1	1			1	3	2	1.50
Aethridae	Hepathus epheliticus	Cangrejo fuente	Ringtraps	3	1	1			1	2	2	1.67
Menippidae	Menippe mercenaria	Cangrejo	Ringtraps	1	1	1			3	3	3	2.00
Epialthidae	Libinia duvia	Maxquíl	Ringtraps	1	1	2			1	2	3	1.67
Gerreidae	Eucinostomus gula	Mojarra	Ringtraps	3	3	3	1	1	1	2		2.00
Gerreidae	Diapterus rhombeus	Mojarra	Ringtraps	3	3	3	1	1	2	2		2.14
Ariidae	Bagre marinus	Bagre 2	Ringtraps	2	2	3	1	1	3	3		2.14
Ariidae	Ariopsis melanopus	Bagre 3	Ringtraps	2	2	3	1	1	3	3		2.14

Family name	Scientific name	Common name	Fishery descriptor	Susceptibility Scores [1-3]					Cumulative only					MSC PSA-derived score	Risk Category Name	MSC scoring guidepost
				Availability	Encounterability	Selectivity	Post-capture mortality	Total (multiplicative)	PSA Score	Catch (tons)	Weighting	Weighted Total	Weighted PSA Score			
Tetraodontidae	Sphoeroides testudineus	Botete	Ringtraps	2	3	2	2	1.58	2.03	10000	1.00	2.03	2.03	95	Low	≥80
Ariidae	Ariopsis felis	Bagre	Ringtraps	1	3	3	3	1.65	2.70	10000	1.00	2.70	2.70	81	Low	≥80
Sparidae	Archosargus rhomboidalis	Postá	Ringtraps	2	3	3	3	2.33	2.73	10000	1.00	2.73	2.73	81	Low	≥80
Achiridae	Achirus lineatus	Lenguado	Ringtraps	1	3	3	3	1.65	2.59	10000	1.00	2.59	2.59	84	Low	≥80
Diogenidae	Clibanarius vittatus	Cangrejo hermitaño	Ringtraps	1	3	2	1	1.13	2.44	10000	1.00	2.44	2.44	87	Low	≥80
Portunidae	Callinectes similis	Jaiba pata seca	Ringtraps	3	3	2	3	2.33	2.77	10000	1.00	2.77	2.77	80	Low	≥80
Portunidae	Callinectes rathbunae	Jaiba negra o café	Ringtraps	1	3	3	3	1.65	2.23	50000	1.00	2.23	2.23	92	Low	≥80
Aethridae	Hepathus epheliticus	Cangrejo fuente	Ringtraps	1	3	3	2	1.43	2.19	50000	1.00	2.19	2.19	92	Low	≥80
Menippidae	Menippe mercenaria	Cangrejo	Ringtraps	2	3	3	1	1.43	2.46	30000	1.00	2.46	2.46	87	Low	≥80
Epialthidae	Libinia duvia	Maxquíl	Ringtraps	1	3	3	3	1.65	2.35	50000	1.00	2.35	2.35	89	Low	≥80
Gerreidae	Eucinostomus gula	Mojarra	Ringtraps	1	3	3	3	1.65	2.59	10000	1.00	2.59	2.59	84	Low	≥80
Gerreidae	Diapterus rhombeus	Mojarra	Ringtraps	1	3	3	3	1.65	2.70	200	1.00	2.70	2.70	81	Low	≥80
Ariidae	Bagre marinus	Bagre 2	Ringtraps	1	3	3	3	1.65	2.70	10000	1.00	2.70	2.70	81	Low	≥80
Ariidae	Ariopsis melanopus	Bagre 3	Ringtraps	1	3	3	3	1.65	2.70	10000	1.00	2.70	2.70	81	Low	≥80



## Conclusions

Compared to other coastal lagoons in the Gulf of Mexico, Laguna de Términos has a high species richness that responds to the environmental heterogeneity in terms of salinity and types of substrate that characterizes this lagoon (Raz-Guzman, 2010). Through the Biological Monitoring Program 14 species were identified as species associated to the blue crab fishery and its fishermen. **The Productivity Susceptibility Analysis results show a low risk score to all of them which allows a score of 90 in the Performance Indicator 2.2.1.** of the MSC standard. Specially attention should be taken to species that are part of other fisheries such as *Menippe mercenaria* and *Libinia duvia*.

*Menippe mercenaria* has a specific fishery with 783 fishing permits (783 traps) in Campeche ([www.pescandodatos.org](http://www.pescandodatos.org), updated August, 2019) none of them in Laguna de Términos or Isla Aguada. Even though, when found it is targeted as of the high value of its chelae. This fishery does not pose a mortality risk since most fishermen only take one chelae, but it is demonstrated this practice diminishes its fecundity.

*Libinia duvia* is used as bait for the octopus's fishery. This fishery takes place from August to mid- December, representing a threat only during this period of the year.

Less information is available on crustaceans.

## Literature cited

**Abascal-Monroy, I.M., M.J. Zetina-Rejón, F. EscobarToledo, G.A. López-Ibarra, A. Sosa-López & A. Tripp-Valdez.** 2016. Functional and structural food web comparison of Términos Lagoon, Mexico in three periods (1980, 1998, and 2011). *Estuar. Coast.*, 39: 1282-1293

**Abeels HA, Loh AN, Volety AK.** 2012. Trophic transfer and habitat use of oyster *Crassostrea virginica* reefs in southwest Florida, identified by stable isotope analysis. *Marine Ecology Progress Series* 462:125-142

**Aceró A.** 2002. Order Siluriformes Ariidae sea catfishes. En: Carpenter K, editores. The living marine resources of the western central Atlantic. Volumen 2. Roma: Food and Agriculture Organization of the United Nations (FAO). p. 831-852.

**Adam E. Cohen, Gary P. Garrett, Melissa J. Casarez, Dean A. Hendrickson, Benjamin J. Labay, Tomislav Urban, John Gentle, Dennis Wylie, and David Walling.** 2018. Conserving Texas Biodiversity: Status, Trends, and Conservation Planning for Fishes of Greatest Conservation Need. Final Report.

**Aguirre I. A., Díaz R.S,** 2000. Estructura poblacional, madurez gonádica y alimentación de *Eugerres plumiere* (gerreidae) en el sistema fluvio-deltaico Aom-Atasta, México. *Ciencias Marinas*, vol. 26, num 2, junio. Pp. 253-273.

**Amador-del Angel, L.E., E. Guevara-Carrió, R. Brito and A.T. Wakida-Kusunoki,** 2015. Length-weight relationships of fish species associated with the mangrove forest in the southwestern Terminos Lagoon, Campeche (Mexico). *J. Appl. Ichthyol.* 31(1):228-230.

**Armstrong, M. P., M. D. Murphy, R. G. Muller, D. P. Harshany, and R. E. Crabtree.** 1996. A stock assessment of Hardhead Catfish, *Arius felis*, and Gafftopsail Catfish, *Bagre marinus*, in Florida waters. Florida Department of Environmental Protection, Florida Marine Research Institute, St. Petersburg

**Ayala-Pérez L.A., J. Ramos M. D., Flores H. A. Sosa L., G. E. Martínez,** 2015. Ictiofauna Marina Costera de Campeche, Instituto Epomex Instituto de Ecología, Pesquerías y Oceanografía del Golfo de México. 502.

**Bonilla Gómez, J. L., Jorge A. López-Rocha, Maribel Badillo Alemán, Juani Tzeek Tuz and Xavier Chiappa-Carrara.** 2011. Growth and Mortality of *Lagodon Rhomboides* (Pisces: Sparidae) In A Tropical Coastal Lagoon In Northwestern Yucatan, Mexico. *Rev. Mar. Cost.* ISSN 1659-455X. Vol. 3: 99-109, Diciembre 2011.

**Bonilla-Gomez, J.L. J. Tzeek-Tuz, M. Badillo & X. Chiappa-Carrara,** 2012. Growth and mortality of *Sphoeroides testudineus* (Tetraodontiformes: Tetraodontidae) in a tropical coastal lagoon in northwestern Yucatán Peninsula. *Revista Mexicana de Biodiversidad* 83: 1237-1240, 2012. DOI: 10.7550/rmb.28616

**Bravo-Calderón, A.; J. López-Rocha y H. Cisneros-Reyes.** 2016. Caracterización de la Pesquería de Jaiba en sisal, Yucatán. Número Especial no. 4. Año 6 no. 11. R. Simposio: Red para el conocimiento de los recursos costeros del sureste. 21 pp.

<https://www.researchgate.net/publication/307560004>





**Breder, C.M. and D.E. Rosen**, 1966. Modes of reproduction in fishes. T.F.H. Publications, Neptune City, New Jersey. 941 p.

**Carla Isobel Elliff; Sérgio Luiz dos Santos Tutui; Marcelo Ricardo Souza<sup>1</sup>; Acácio Ribeiro Gomes Tomas**. 2013 Population structure of caitipa mojarra (*Diapterus rhombeus*) in an Estuarine system of southeastern Brazil\* Bol. Inst. Pesca, São Paulo, 39(4): 411 – 421, 2013

**Carmona-Osalde, C. & M. Rodríguez-Serna**, 2012. Reproductive aspects of the spider crab *Libinia dubia* under laboratory conditions. Hidrobiológica vol.22 no.1 México ene./abr. 2012

**Cervigón, F., R. Cipriani, W. Fischer, L. Garibaldi, M. Hendrickx, A.J. Lemus, R. Márquez, J.M. Poutiers, G. Robaina and B. Rodriguez**, 1992. Fichas FAO de identificación de especies para los fines de la pesca. Guía de campo de las especies comerciales marinas y de aguas salobres de la costa septentrional de Sur América. FAO, Rome. 513 p. Preparado con el financiamiento de la Comisión de Comunidades Europeas y de NORAD.

**Cervigón, F.**, 1993. Los peces marinos de Venezuela. Volume 2. Fundación Científica Los Roques, Caracas, Venezuela. 497 p.

**Chavance, P., A. Yañez-Arancibia, D. Flores-Hernández, A.L. Lara-Domínguez and F.A. Linares**, 1986. Ecology, biology and population dynamics of *Archosargus rhomboidalis* (Pisces, Sparidae) in a tropical coastal lagoon system, southern Gulf of Mexico. An. Inst. Cienc. Mar Limnol. Univ. Nac. Auton. Mex. 13(2):11-30.

**Chi-Espínola, A.A., and Vega-Cendejas, M.E.** 2013. Feeding habits of *Sphoeroides testudineus* (Perciformes: Tetraodontidae) in the lagoon system of Ria Lagartos, Yucatan, Mexico. Revista de biología tropical 61(2): 849-858.

**Claudia Carmona-Osalde and Miguel Rodríguez-Serna**, 2012 Reproductive aspects of the spider crab *Libinia dubia* under laboratory conditions Hidrobiológica vol.22 no.1 México ene./abr. 2012 versión impresa ISSN 0188-8897

**Cohen, Adam E., Gary P. Garrett, Melissa J. Casarez, Dean A. Hendrickson, Benjamin J. Labay, Tomislav Urban, John Gentle, Dennis Wiley, and David Walling**. 2018. "Final Report: 'Conserving Texas Biodiversity: Status, Trends, and Conservation Planning for Fishes of Greatest Conservation Need' (Contract No. 459125 UTA14-001402)," Texas Parks and Wildlife Department through U.S. Fish and Wildlife Service State Wildlife Grant Program, grant TX T-106-1 (CFDA# 15.634), January, 362 pages. <https://doi.org/10.15781/T26M33M7Z>

**Comisión Nacional para el Conocimiento y Uso de la Biodiversidad (CONABIO)**. 2010. Ficha técnica para la evaluación de los sitios prioritarios para la conservación de los ambientes costeros y oceánicos de México. Humedales costeros y Plataforma continental de Cabo Catoche. Mesa de Trabajo: Golfo de México (Ecorregión Nivel-I CCA: Golfo de México Sur; Categoría del sitio: Zona costera). 7 p.

**Crowley CE, Gandy RL, Daly KL, Leone EH**. 2018. Assessment of Maturity in the Eastern Gulf of Mexico Florida Stone Crab *Menippe mercenaria* Fishery. Transactions of the American Fisheries Society 147:1124–1145

**Crowley, C.E., C. P. Shea, R. L. Gandy, K. Daly.** 2019. Fecundity assessment of Stone Crabs in the Eastern Gulf of Mexico. *Marine & Coastal Fisheries. Dynamics, Management, and Ecosystem Science*. Vol 11, Issue 1. Pages 32-47. <https://doi.org/10.1002/mcf2.10059>

**de Lima PA, Bertini G, Fransozo V, Gregati RA, Fernandes-Góes LC, Castilho AL.** 2014. Reproductive biology of *Hepatus pudibundus* (Crustacea: Brachyura), the most abundant crab on the southeastern Brazilian coast. *Biologia* 69(2):219-227

**Dragovich, AJ & JA Kelly, Jr.** 1964. Ecological observations of macro-invertebrates in Tampa Bay, Florida. *Bull. Mar. Sci.* 14: 74-102.

**Enzenross, R & L Enzenross.** 2000. Non-Mediterranean crustaceans in Tunisian waters (Decapoda, Macrura and Brachyura). *Crustaceana* 73: 187-195.

**Felder, D.L., F. Alvarez, J.W. Goy and R. Lemaitre.** 2009 Decapoda (Crustacea) of the Gulf of Mexico, with comments on the Amphionidacea. pp. 1019-1104 In Felder, D.L.; Camp, D.K. (eds.) *Gulf of Mexico - Origins, Waters, and Biota. Biodiversity*. Texas A&M University Press, College Station, Texas.

**Fernandes-Miazaki, L.; S. Morilhas-Simoes; A. Leao- Castilho and R. Caetano-Costa.** 2018. Population dynamics of the crab *Hepatus pudibundus* (Herbst, 1785) (Decapoda, Aethridae) on the southern coast of Sao Paulo state, Brazil. *Journal of Marine Biological Association of the United Kingdom*. 1–12. <https://doi.org/10.1017/S0025315418000620>

**Flinn S., Midway S, A. Ostrowski.** 2019. Age and Growth of Hardhead catfish and gafftopsail catfish in coastal Louisiana, USA. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* 11:362–371, 2019. DOI: 10.1002/mcf2.10089

**Franks, J.S., J.Y. Christmas, W.L. Siler, R. Combs, R. Waller & C. Burns.** 1972. A study of nektonic and benthic faunas of the shallow gulf of Mexico off the state of Mississippi as related to some physical, chemical and geological factors. *Gulf Res. Rep.*, 4: 1- 148.

**Froese, R. and D. Pauly.** Editors. 2019. FishBase. World Wide Web electronic publication.

**Froese, R., N. Demirel, G. Coro, K.M. Kleisner and H. Winker,** 2017. Estimating fisheries reference points from catch and resilience. *Fish and Fisheries* 18(3):506-526.

**Gandy R, Crowley C, Chagris D, Crawford C.** 2016. The effect of temperature on release mortality of declawed *Menippe mercenaria* in the Florida stone crab fishery. *Bull Mar Sci* 92(1):1-15

**Gandy RL, Crowley CE, Leone EH, Crawford CR.** 2018. Increasing the Selectivity of the Stone Crab *Menippe mercenaria* Trap by the Addition of a Cull Ring. *North American Journal of Fisheries Management* 38:1275–1283

**García-Montes, J.F.; A. Gracia; L. A. Soto.** 1987. Morfometría, crecimiento relativo y fecundidad de la jaiba del Golfo, *Callinectes similis* Williams, 1966 (Decapoda: Portunidae). Morphometry, Relative Growth and Fecundity of the Gulf Crab, *Callinectes similis* Williams, 1966. *Ciencias Marinas* 13(4):137-161, 1987.

**Gerhart DS and Bert TM.** 2008. Life-history aspects of stone crabs (Genus *Menippe*): Size at maturity, growth, and age. *Journal of Crustacean Biology* 28(2):252-261



**Geslaine Rafaela Lemos Gonçalves, Lizandra Fernandes Miazaki, Eduardo Antonio Bolla Junior, Raphael Cezar Grabowski, Rogério Caetano da Costa, Antonio Leão Castilho**, 2020. Growth and longevity of the spider crab *Libinia ferreirae* (Majoidea, Epialtidae) Nauplius, 28: e2020003. DOI 10.1590/2358-2936e2020003

**Gines, H. and F. Cervigón**, 1967. Exploración pesquera en las costas de Guyana y Surinam año 1967. Estación de Investigaciones Marinas de Margarita. Fundación La Salle de Ciencias Naturales, no. 29.

**Gulf of Mexico Fishery Management Council**. 1979 . (GMFMC) Fishery management plan for the stone crab fishery in the Gulf of Mexico Federal Register 44: 19445–19496.

**Hernández, P.; A. Rombenso, M. A.A. Pinheiro & Nuno Simoes**. 2012. Population structure and sexual maturity of the calico box crab *Hepatus epheliticus* Linnaeus (Brachyura, Hepatidae) from Yucatan Peninsula, Mexico. Lat. Am. J. Aquat. Res., 50(2): 480-486. DOI: 10.3856/vol40-issue2-fulltext-25

**J. Labay, Tomislav Urban, John Gentle, Dennis Wiley, and David Walling**. 2018. “Final Report: ‘Conserving Texas Biodiversity: Status, Trends, and Conservation Planning for Fishes of Greatest Conservation Need’ (Contract No. 459125 UTA14-001402),” Texas Parks and Wildlife Department through U.S. Fish and Wildlife Service State Wildlife Grant Program, grant TX T-106-1 (CFDA# 15.634), January, 362 pages. <https://doi.org/10.15781/T26M33M7Z>

**J. Sánchez and A. Raz-Guzman**, 1997. Distribution patterns of tropical estuarine brachyuran crabs in the Gulf of Mexico. Journal of crustacean biology, 17(4): 609-620, 1997

**Keith, P., P.-Y. Le Bail and P. Planquette**, 2000. Atlas des poissons d'eau douce de Guyane. Tome 2, Fascicule I: Batrachoidiformes, Mugiliformes, Beloniformes, Cyprinodontiformes, Synbranchiformes, Perciformes, Pleuronectiformes, Tetraodontiformes. Collection Patrimoines Naturels 43(I): 286p. Paris: Publications scientifiques du Muséum national d'Histoire naturelle. DOI 2-86515-117-4

**Keunecke KA, D'Incao F, Fonseca DB**. 2007 . Growth and mortality of *Hepatus pudibundus* (Crustacea: Calappidae) in south-western Brazil. Journal of the Marine Biological Association of the United Kingdom 87:885- 891

**Lieske, E. and R. Myers**, 1994. Collins Pocket Guide. Coral reef fishes. Indo-Pacific & Caribbean including the Red Sea. Haper Collins Publishers, 400 p. DOI 0-00-219974-2

**Mallard-Colmenero, L., A. Yáñez-Arancibia & F. Amezcua-Linares**. 1982. Taxonomía, diversidad, distribución y abundancia de los tetraodontidos de la Laguna de Términos, sur del Golfo de México. (Pisces: Tetraodontidae). An. Inst. Cienc. del Mar y Limnol. Univ. Nal. Autón. Mex. 9: 161-212.

**Mantelatto FLM and Patracco M**. 1997. Natural diet of the crab *Hepatus pundibundus* (Brachyura: Calappidae) in Fortaleza Bay, Ubatuba (SP), Brazil. Journal of Crustacean Biology 17(3):440-446

**Mascaro, M., A.M. Castillo, N. Simoes and X. Chiappa-Carrara** 2007 Variations in the feeding habits of *Callinectes rathbunae* in Las Palmas lagoon (southern Gulf of Mexico) on three temporal scales. Crustaceana 80(2):139-160.



**Mascaro, M., A.M. Castillo, N. Simoes and X. Chiappa-Carrara** 2007 Variations in the feeding habits of *Callinectes rathbunae* in Las Palmas lagoon (southern Gulf of Mexico) on three temporal scales. *Crustaceana* 80(2):139-160.

**Mexicano-Cíntora, G.**, 1999. Crecimiento y reproducción de la mojarra, *Eucinostomus gula* de Celestún, Yucatán, México. *Proc. Gulf Carribb. Fish. Inst.* 45:524-536.

**Morales-Zárata, M.V.; F. Arreguín-Sánchez; J. López-Martínez; S.E. Lluch-Cota.** 2004. Ecosystem trophic structure and energy flux in the Northern gulf of California, México. *Ecological Modelling* 174 (2004) 331-345

**Nedgreiros-Fransozo, A. Fransozo & G. L. Hirose.** 2008. The megalopa and early juvenile development of *Hepatus pudibundus* (Crustacea: Brachyura: Aethroldea) reared from neuston samples. *Revista Brasileira de Zoología* 25(4): 608-616, December, 2008.

**Patricio Hernández, Artur Rombenso , Marcelo A.A. Pinheiro & Nuno Simões.** 2012. Population structure and sexual maturity of the calico box crab *Hepatus epheliticus* Linnaeus (Brachyura, Hepatidae) from Yucatan Peninsula, Mexico *Lat. Am. J. Aquat. Res.*, 40(2): 480-486, 2012

**Pauly, D.**, 1991. Growth of the checkered puffer *Sphoeroides testudineus*: postscript to papers by Targett and Pauly & Ingles. *Fishbyte* 9(1):19-22.

**Pech, D.; S. B. Balam; A. León H.; E. Nuñez-Lara y P. Rodríguez-Pliego.** 2015. Los macro invertebrados bentónicos de laguna de Términos: ¿Cuántos son y cómo se distribuyen?. 167-180 pp. En: Ramos Miranda J., y G.J. Villalobos Zapata (editores), 2015. Aspectos socioambientales de la región de la laguna de Términos, Campeche. Universidad Autónoma de Campeche. 210 p.

**Pedro Arenas-Granados I y Arturo Acero P.** 1992. Organización trófica de las mojarra (Pisces: Gerreidae) de la Ciénaga Grande de Santa Marta (Caribe Colombiano) *Rev. Biol. Trop.*, 40(3): 287-302, 1992

**Peniche-Pérez J.C., González-Salas C., Villegas-Hernández H., Díaz-Gamboa R., Aguilar-Perera A., GuillenHernández S., Poot-López G.R.** 2019. Reproductive biology of the southern pufferfish, *Sphoeroides nephelus* (Actinopterygii: Tetraodontiformes: Tetraodontidae), in the northern coast off the Yucatan Peninsula, Mexico. *Acta Ichthyol. Piscat.* 49 (2): 133-146.

**Piller SC, Henry RP, Doeller JE and DW Kraus.** 1995. A comparison of the gill physiology of two euryhaline crab species, *Callinectes sapidus* and *Callinectes similis*: energy production, transport-related enzymes and osmoregulation as a function of acclimation salinity. *Journal of Experimental Biology* 198:349-358.

**Raz-Guzmán, A., A.J. Sánchez, L.A. Soto, y F. Alvarez,** 1986. Catálogo ilustrado de cangrejos braquiuros y anomuros de laguna de Términos, Campeche (Crustacea: Brachyura, Anomura). *Anales del Instituto de Biología, Serie Zoología*, 57(2): 343-383.

**Raz-Guzman,**2010. Estudio de caso: crustáceos de la laguna de Términos. En. *Diversidad de especies: Estudio de Caso.* 275-278 pp.



**Reigada, ALD; Negreirosfransozo, M. L.** 1995 Fecundity of crab Hepatus-pudibundas (herbst, 1785) (crustacea, decapoda, calappidae) in Ubatuba (SP), Brazil. Arquivos de Biologia E Tecnologia. Curitiba-Paraná: Inst Tecnologia Parana, v. 38, n. 2, p. 661-668, 1995. <<http://hdl.handle.net/11449/31537>>.

**Reis-Filho J. A. O.H.C. Leduc,** 2018. Mysterious and elaborated: The reproductive behavior of the rhomboid mojarra, *Diapterus rhombeus* (Cuvier, 1829) in Brazilian mangrove habitats. Helgoland Marine Research s (2018) 72:7

**Rodrigues da Costa, Marcus; Rafael Jardim Albieri; Leonardo Mitrano Neves; Alex Braz Lacone Santos; Francisco Gerson Araujo.** 2012. Distribution and size of the mojarra *Diapterus rhombeus* (Cuvier) (Actinopterygii, Gerreidae) in a Southeastern Brazilian bay. In: Brazilian Journal of Oceanography On-line version ISSN 1982-436X Braz. j. oceanogr. vol. 60 no.2 São Paulo Apr./June 2012 <https://doi.org/10.1590/S1679-87592012000200010>

**Robins, C.R. and G.C. Ray,** 1986. A field guide to Atlantic coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 354 p.

**Ruppert, E.E., R.S. Fox and R.D. Barnes** 2004 Invertebrate Zoology. A functional evolutionary approach. 7th Ed. Brooks/Cole, Thomson Learning learning, Inc. 990 p.

**Seafood Watch Consulting Researcher,** 2019; Seafood Watch Stone crab United States of America: Gulf of Mexico & Western Central Atlantic Pots. Evaluation. 33pp.

**Segura-Berttolini, E.C.** 2011, Biología reproductiva de los machos de Bagre marinus (PISCES: ARIIDAE), en el Sureste del Golfo de México. Tesis de Maestría. El Colegio de la Frontera Sur (ECOSUR), 111 pp.

**Sergio Chazaro-Olvera; Arturo Rocha-Ramirez; Ramiro Roman-Contreras** (2000). Observations on feeding, maturity and fecundity of *Callinectes similis* Williams, 1966, on the central continental shelf off Veracruz, Gulf of Mexico". *Crustaceana*. 73 (3): 323-332. doi:10.1163/156854000504417

**Shao, K., Matsuura, K., Leis, J.L., Hardy, G., Larson, H. & Liu, M.** 2014. *Sphoeroides testudineus*. The IUCN Red List of Threatened Species 2014: e.T193813A2281154. <https://dx.doi.org/10.2305/IUCN.UK.2014-3.RLTS.T193813A2281154.en>. Downloaded on 08 September 2020.

**Soto, L. A.,** 1980. Decapod crustacea shelf-fauna of the Campeche Bank: Fishery and ecological aspects. *Gulf Caribb. Fish. Inst.* 32: 66-81.

**Smith, A.D.M., Fulton, E.J., Hobday, A.J., Smith, D.C. & Shoulder, P.** 2007. Scientific tools to support the practical implementation of ecosystem-based fisheries management. *ICES Journal of Marine Science*, 64: 633-639.

**Tavares, M.** 2003 True Crabs. pp. 327-352. In Carpenter, K.E. (ed.) The living marine resources of the Western Central Atlantic. Volume1: introduction, molluscs, crustaceans, hagfishes, sharks, batoid fishes, and chimaeras. FAO Species Identification Guide for Fishery Purposes and American Society of Ichthyologists and Herpetologists Special Publication No. 5. Rome, FAO. pp. 1-600. Pp 341.



**Tabb, DC & RB Manning.** 1961. A checklist of the flora and fauna of northern Florida Bay and adjacent brackish waters of the Florida mainland collected during the period July, 1957 through September, 1960. *Bull. Mar. Sci.* 11: 552-649.

**Tunberg, BG & SA Reed.** 2004. Mass occurrence of the jellyfish *Stomolophus meleagris* and an associated spider crab *Libinia dubia*, eastern Florida. *FL Scientist* 67: 93-104.

**Vega-Cendejas, M.E. and Hernández de Santillana, M.** 2004. Fish community structure and dynamics in a coastal hypersaline lagoon: Rio Lagartos, Yucatan, Mexico. *Estuarine Coastal and Shelf Science* 60(2): 285-299.

**Williams, A.B.** 1984. *Shrimps, Lobsters, and Crabs of the Atlantic Coast of the Eastern United States, Maine to Florida.* Washington, D.C.: Smithsonian Institution Press, 550 pp.

**Winemiller, K. O., and K. A. Rose.** 1992. Patterns of life-history diversification in North American fishes: implications for population regulation. *Canadian Journal of Fisheries and Aquatic Sciences* 49:2196-2218.

<https://animaldiversity.org>

<https://www.crabdatabase.info/en/crabs>

<https://www.fishbase.de>

<https://naturalhistory2.si.edu>

<https://www.sealifebase.se>